

- [1] Anderson LE, Inglehart MR, El-Kholy K, Eber R, Wang HL: Implant associated soft tissue defects in anterior maxilla: a randomized control trial comparing subepithelial connective tissue graft and acellular dermal matrix allograft.: *Implant Dent* (4),416–25 (2014).
- [2] Araujo MG, Lindhe J: Dimensional ridge alterations following tooth extraction. An experimental study in the dog. *J Clin Periodontol* 32, 212–218 (2005).
- [3] Araujo M, Linder E, Wennström J, Lindhe J: The Influence of Bio-Oss Collagen on Healing of an Extraction Socket: An Experimental Study in the Dog. *Int J Periodontics Restorative Dent* 28,123–135 (2008).
- [4] Basetti RG, Stähli A, Basetti MA, Scuean A: Soft tissue augmentation procedures at second- stage surgery: a systematic review. *Clin Oral Investig* 2016 Apr. 4. (Epub ahead of print).
- [5] Block MS et al.: Hydroxyapatite-coated cylindrical implants in the posterior mandible: 10-year observations. *Int J Oral Maxillofac Implants* 11, 626–633 (1996).
- [6] Cardaropoli D, Cardaropoli G: Preservation of the postextraction alveolar ridge: a clinical and histologic study. *Int J Periodontics Restorative Dent* 28 (5), 469–477 (2008).
- [7] Chiu YW, Lee SY, Lin YC, Lai YL: Significance of the width of keratinized mucosa on peri-implant health. *J Chin Med Assoc* 78 (7), 389–94 (2015).
- [8] Deporter DA et al: A histological evaluation of a functional endosseous, porous-surfaced, titanium alloy dental implant system in the dog. *J Dent Res* 67, 1190–1195 (1988).
- [9] Fickl S: Peri-implant mucosal recession: Clinical significance and therapeutic opportunities. *Quintessence Int* 46 (8), 671–6 (2015).
- [10] Kirsch A, Ackermann KL: The IMZ osteointegrated implant system. *Dent Clin North Am* 33, 733–791 (1989).
- [11] Lee JH, Lee JS, Baek WS, Lim HC, Cha JK, Choi SH, Jung UW: Assessment of dehydrothermally cross-linked collagen membrane for guided bone regeneration around peri-implant dehiscence defects: a randomized single-blinded clinical trial. *J Periodontal Implant Sci* 45 (6), 229–37 (2015).
- [12] Lima RS, Peruzzo DC, Napimoga MH, Saba-Chuifi E, Santos-Pereira SA, Martinez EF: Evaluation of the Biological Behavior of Mucograft in Human Gingival Fibroblasts: An In Vitro Study. *Braz Dent J* 26 (6), 602–6 (2015).
- [13] Lin GH, Chan HL, Wang HLJ: The significance of keratinized mucosa on implant health: a systematic review. *J Periodontol* 84 (12), 1755–67 (2013).
- [14] Liu C, Su Y, Tan B, Ma P, Wu G, Li J, Geng W: Reconstruction of attached soft tissue around dental implants by acellular dermal matrix grafts and resin splint. *Int J Clin Exp Med* 7 (12), 4666–76 (2014).
- [15] McGuire MK, Scheyer ET: Xenogeneic collagen matrix with coronally advanced flap compared to connective tissue with coronally advanced flap for the treatment of dehiscence-type recession defects. *J Periodontol* 81 (8), 1108-17 (2010).
- [16] Roos-Jansaker AM et al: Nine-to fourteen-year follow-up of implant treatment. Part I: implant loss and associations to various factors. *J Clin Periodontol* 33, 283–289 (2006).
- [17] Roos-Jansaker AM et al: Nine-to fourteen-year follow-up of implant treatment. Part II: presence of peri-implant lesions. *J Clin Periodontol* 33, 290–295 (2006).
- [18] Roos-Jansaker AM et al: Nine-to fourteen-year follow-up of implant treatment. Part I: factors associated with peri-implant lesions. *J Clin Periodontol* 33, 296–301 (2006).

- [19] Sanz M et al: Clinical evaluation of a new collagen matrix (Mucograft prototype) to enhance the width of keratinized tissue in patients with fixed prosthetic restorations: a randomized prospective clinical trial. *J Clin Periodontol* 36 (10), 868–76 (2009).
- [20] Schmitt CM, Moest T, Lutz R, Werhahn F, Neukam FW, Schlegel KA: Long-term outcomes after vestibuloplasty with a porcine collagen matrix (Mucograft) versus the free gingival graft: a comparative prospective clinical trial. *Clin Oral Implants res.* 2015 Feb 27. (Epub ahead of print).
- [21] Schropp L, Wenzel A, Kostopoulos L, Karring T: Bone healing soft tissue contour changes following single-tooth extraction: A clinical and radiographic 12-month prospective study. *Int J Periodontics Restorative Dent* 23, 313–323 (2003).
- [22] Strietzel FP, Shakibaie MB: Der Einsatz der TefGen-FD®-Membran zum Erhalt des Alveolarkamms nach Zahnextraktionen: eine klinische Studie. *Dtsch Zahnärztl Z* 12, 883–886 (1998).

Forcierte Extrusion – ein Behandlungskonzept für bisher ausweglose Fälle: Ein biologisches Konzept zum Erhalt tief zerstörter bzw. frakturierter Zähne

- [1] Baumgart R, Kuhn V, Hinterwimmer S, Krammer M, Mutschler Wolf: Zugkraftmessungen beim knöchernen Segmenttransport – in vivo Untersuchungen am Menschen. *Biomedizinische Technik* 49 (9), 248–256 (2004).
- [2] Berglundh T, Marinello CP, Lindhe J, Thilander B, Liljenberg B: Periodontal tissue reactions to orthodontic extrusion. An experimental study in the dog. *J Clin Periodontol* 18 (5), 330–336 (1991).
- [3] Berglundh T, Lindhe J: Dimension of the periimplant mucosa. Biological width revisited. *J Clin Periodontol* 23 (10), 971–973 (1996).
- [4] Bondemark L, Kurol J, Hallonsten AL, Andreasen JO: Attractive magnets for orthodontic extrusion of crown-root fractured teeth. *Am J Orthod Dentofacial Orthop* 112 (2), 187–193 (1997).
- [5] Carvalho CV, Bauer FP, Romito GA, Pannuti CM, De Micheli G: Orthodontic extrusion with or without circumferential supracrestal fiberotomy and root planing. *Int J Periodontics Restorative Dent* 26 (1), 87–93 (2006).
- [6] Cooke MS, Scheer B: Extrusion of fractured teeth. The evolution of practical clinical techniques. *Br Dent J* 149 (2), 50–53 (1980).
- [7] Coval NM: Physiological orthodontic extrusion of fractured incisors. *Chronicle* 36 (10), 252–256 (1973).
- [8] Gargiulo AW, Wentz FM, Orban B: Dimensions and relations of the dentogingival junction in humans. *J Periodontol* 32, 261–267 (1961).
- [9] Hamilton RS, Gutmann JL: Endodontic-orthodontic relationships: a review of integrated treatment planning challenges. *Int Endod J* 32 (5), 343–360 (1999).
- [10] Han G, Huang S, Von den Hoff JW, Zeng X, Kuijpers-Jagtman AM: Root resorption after orthodontic intrusion and extrusion: an intraindividual study. *Angle Orthod* 75 (6), 912–918 (2005).
- [11] Harder S, Mehl C, Kern M: Behandlungsmöglichkeiten bei tief frakturen Pfeilerzähnen. *Quintessenz* 61, 1485–1494 (2010).
- [12] Ingber JS: Forced eruption. Part I. *J Periodontol* 45, 199–206 (1974).
- [13] Ingber JS: Forced eruption: part II. A method of treating nonrestorable teeth. Periodontal and restorative considerations. *J Periodontol* 47 (4), 203–216 (1976).
- [14] Johnston CD, Littlewood SJ: Retention in orthodontics. *Br Dent J* 218 (3), 119–122 (2015).
- [15] Kim SH, Tramontina VA, Papalexiou V, Luczyszyn SM, Grassi MB, de Fatima Scarpim M, Tanaka OM: Rapid orthodontic extrusion using an interocclusal appliance for the reestablishment of biologic width: a case report. *Quintessence Int* 42 (3), 201–204 (2011).
- [16] Kina JR, Dos Santos PH, Kina EF, Suzuki TY, Dos Santos PL: Periodontal and prosthetic biologic considerations to restore biological width in posterior teeth. *J Craniofac Surg* 22 (5), 1913–1916 (2011).
- [17] Korayem M, Flores-Mir C, Nassar U, Olfert K: Implant site development by orthodontic extrusion. A systematic review. *Angle Orthod* 78 (4), 752–760 (2008).
- [18] Lanning SK, Waldrop TC, Gunsolley JC, Maynard JG: Surgical crown lengthening: evaluation of the biological width. *J Periodontol* 74 (4), 468–474 (2003).
- [19] Lindhe J, Nyman S: Alterations of the position of the marginal soft tissue following periodontal surgery. *J Clin Periodontol* 7 (6), 525–530 (1980).
- [20] Lovdahl PE: Periodontal management and root extrusion of traumatized teeth. *Dent Clin North Am* 39 (1), 169–179 (1995).

- [21] Matthews D: Conclusive support for mechanical nonsurgical pocket therapy in the treatment of periodontal disease. How effective is mechanical nonsurgical pocket therapy? *Evid Based Dent* 6 (3), 68–69 (2005).
- [22] Mehl C, Wolfart S, Kern M: Orthodontic extrusion with magnets: a case report. *Quintessence Int* 39 (5), 371–379 (2008).
- [23] Neumeyer S, Wachtel H: Die Replantation und orthodontische Extrusion hoch resezierter Zähne. *Quintessenz* 60, 1141–1149 (2009).
- [24] Neumeyer S: yah-. Masterthese MOP Freiburg 2010.
- [25] Persson M, Serneke D: Orthodontic extrusion of tooth with cervical root fracture facilitating crown preparation. *Tandlakartidningen* 69 (22), 1263–1269 (1977).
- [26] Rasperini G, Siciliano VI, Cafiero C, Salvi GE, Blasi A, Aglietta M: Crestal bone changes at teeth and implants in periodontally healthy and periodontally compromised patients. A 10-year comparative case-series study. *J Periodontol* 85 (6), e152–159 (2014).
- [27] Rupp R: Root resorption related to orthodontics and other factors: a review of the literature. *J Gen Orthod* 6 (3), 25–29 (1995).
- [28] Simon JH, Lythgoe JB, Torabinejad M: Clinical and histologic evaluation of extruded endodontically treated teeth in dogs. *Oral Surg Oral Med Oral Pathol* 50 (4), 361–371 (1980).
- [29] Tarnow DP, Magner AW, Fletcher P: The effect of the distance from the contact point to the crest of bone on the presence or absence of the interproximal dental papilla. *J Periodontol* 63 (12), 995–996 (1992).
- [30] Whitworth JM, Walls AW, Wassell RW: Crowns and extra-coronal restorations: endodontic considerations: the pulp, the root-treated tooth and the crown. *Br Dent J* 192 (6), 315–320, 323–327 (2002).
- [31] Will LA: Stability and retention. *Front Oral Biol* 18, 56–63 (2016).
- [32] Young L, Binderman I, Yaffe A, Beni L, Vardimon AD: Fiberotomy enhances orthodontic tooth movement and diminishes relapse in a rat model. *Orthod Craniofac Res* 16 (3), 161–168 (2013).

- [1] Anderson LE, Inglehart MR, El-Kholy K, Eber R, Wang HL: Implant associated soft tissue defects in anterior maxilla: a randomized control trial comparing subepithelial connective tissue graft and acellular dermal matrix allograft.: *Implant Dent* (4),416–25 (2014).
- [2] Araujo MG, Lindhe J: Dimensional ridge alterations following tooth extraction. An experimental study in the dog. *J Clin Periodontol* 32, 212–218 (2005).
- [3] Araujo M, Linder E, Wennström J, Lindhe J: The Influence of Bio-Oss Collagen on Healing of an Extraction Socket: An Experimental Study in the Dog. *Int J Periodontics Restorative Dent* 28,123–135 (2008).
- [4] Basetti RG, Stähli A, Basetti MA, Scuean A: Soft tissue augmentation procedures at second- stage surgery: a systematic review. *Clin Oral Investig* 2016 Apr. 4. (Epub ahead of print).
- [5] Block MS et al.: Hydroxyapatite-coated cylindrical implants in the posterior mandible: 10-year observations. *Int J Oral Maxillofac Implants* 11, 626–633 (1996).
- [6] Cardaropoli D, Cardaropoli G: Preservation of the postextraction alveolar ridge: a clinical and histologic study. *Int J Periodontics Restorative Dent* 28 (5), 469–477 (2008).
- [7] Chiu YW, Lee SY, Lin YC, Lai YL: Significance of the width of keratinized mucosa on peri-implant health. *J Chin Med Assoc* 78 (7), 389–94 (2015).
- [8] Deporter DA et al: A histological evaluation of a functional endosseous, porous-surfaced, titanium alloy dental implant system in the dog. *J Dent Res* 67, 1190–1195 (1988).
- [9] Fickl S: Peri-implant mucosal recession: Clinical significance and therapeutic opportunities. *Quintessence Int* 46 (8), 671–6 (2015).
- [10] Kirsch A, Ackermann KL: The IMZ osteointegrated implant system. *Dent Clin North Am* 33, 733–791 (1989).
- [11] Lee JH, Lee JS, Baek WS, Lim HC, Cha JK, Choi SH, Jung UW: Assessment of dehydrothermally cross-linked collagen membrane for guided bone regeneration around peri-implant dehiscence defects: a randomized single-blinded clinical trial. *J Periodontal Implant Sci* 45 (6), 229–37 (2015).
- [12] Lima RS, Peruzzo DC, Napimoga MH, Saba-Chuifi E, Santos-Pereira SA, Martinez EF: Evaluation of the Biological Behavior of Mucograft in Human Gingival Fibroblasts: An In Vitro Study. *Braz Dent J* 26 (6), 602–6 (2015).
- [13] Lin GH, Chan HL, Wang HLJ: The significance of keratinized mucosa on implant health: a systematic review. *J Periodontol* 84 (12), 1755–67 (2013).
- [14] Liu C, Su Y, Tan B, Ma P, Wu G, Li J, Geng W: Reconstruction of attached soft tissue around dental implants by acellular dermal matrix grafts and resin splint. *Int J Clin Exp Med* 7 (12), 4666–76 (2014).
- [15] McGuire MK, Scheyer ET: Xenogeneic collagen matrix with coronally advanced flap compared to connective tissue with coronally advanced flap for the treatment of dehiscence-type recession defects. *J Periodontol* 81 (8), 1108–17 (2010).
- [16] Roos-Jansaker AM et al: Nine-to fourteen-year follow-up of implant treatment. Part I: implant loss and associations to various factors. *J Clin Periodontol* 33, 283–289 (2006).
- [17] Roos-Jansaker AM et al: Nine-to fourteen-year follow-up of implant treatment. Part II: presence of peri-implant lesions. *J Clin Periodontol* 33, 290–295 (2006).
- [18] Roos-Jansaker AM et al: Nine-to fourteen-year follow-up of implant treatment. Part I: factors associated with peri-implant lesions. *J Clin Periodontol* 33, 296–301 (2006).

- [19] Sanz M et al: Clinical evaluation of a new collagen matrix (Mucograft prototype) to enhance the width of keratinized tissue in patients with fixed prosthetic restorations: a randomized prospective clinical trial. *J Clin Periodontol* 36 (10), 868–76 (2009).
- [20] Schmitt CM, Moest T, Lutz R, Werhahn F, Neukam FW, Schlegel KA: Long-term outcomes after vestibuloplasty with a porcine collagen matrix (Mucograft) versus the free gingival graft: a comparative prospective clinical trial. *Clin Oral Implants res.* 2015 Feb 27. (Epub ahead of print).
- [21] Schropp L, Wenzel A, Kostopoulos L, Karring T: Bone healing soft tissue contour changes following single-tooth extraction: A clinical and radiographic 12-month prospective study. *Int J Periodontics Restorative Dent* 23, 313–323 (2003).
- [22] Strietzel FP, Shakibaie MB: Der Einsatz der TefGen-FD[®]-Membran zum Erhalt des Alveolarkamms nach Zahnextraktionen: eine klinische Studie. *Dtsch Zahnärztl Z* 12, 883–886 (1998).

- [1] Andreiotelli M, Att W, Strub JR. Prosthodontic complications with implant overdentures: a systematic literature review. *Int J Prosthodont* 2010; 23:195–203.
- [2] Chaar MS, Att W, Strub JR. Prosthetic outcome of cement-retained implant-supported fixed dental restorations: a systematic review. *J Oral Rehabil* 2011; 38:697–711.
- [3] Hämmerle CH, Wagner D, Brägger U, Lussi A, Karayiannis A, Joss A, Lang NP. Threshold of tactile sensitivity perceived with dental endosseous implants and natural teeth. *Clin. Oral Implants Res* 1995;6:83–90.
- [4] Kinsel RP, Lin D. Retrospective analysis of porcelain failures of metal ceramic crowns and fixed partial dentures supported by 729 implants in 152 patients: patient-specific and implant-specific predictors of ceramic failure. *J Prosthet Dent* 2009;101: 388–94.
- [5] Konstantinidis IK, Kotsakis GA, Gerdes S, Walter MH. Cross-sectional study on the prevalence and risk indicators of periimplant diseases. *Eur J Oral Implantol*. 2015;8:75–88.
- [6] Lewis MB, Klineberg I. Prosthodontic considerations designed to optimize outcomes for single-tooth implants. A review of the literature. *Aust Dent J* 2011; 56:181–192.
- [7] Marxkors R, Wolowski A. Unklare Kiefer-Gesichtsbeschwerden: Abgrenzung zahnärztlich-somatischer von psychischen Ursachen. München; Wien: Hanser, 1999.
- [8] Pjetursson BE, Lang NP. Prosthetic treatment planning on the basis of scientific evidence. *J Oral Rehabil* 2008; 35 Suppl 1:72–9.
- [9] Romeo E, Storelli S. Systematic review of the survival rate and the biological, technical, and aesthetic complications of fixed dental prostheses with cantilevers on implants reported in longitudinal studies with a mean of 5 years follow-up. *Clin Oral Implants Res*. 2012; 23 Suppl 6:39–49.
- [10] Salvi GE, Brägger U. Mechanical and technical risks in implant therapy. *Int J Oral Maxillofac Implants* 2009; 24 (suppl):69–85.

- [1] Arima T, Tomonaga A, Toyota M, Inoue SI, Ohata N, Svensson P (2012) Does restriction of mandibular movements during sleep influence jaw-muscle activity? *J Oral Rehabil* 39 (7): 545–551.
- [2] Bader G, Lavigne G (2000) Sleep bruxism; an overview of an oromandibular sleep movement disorder. REVIEW ARTICLE. *Sleep Med Rev* 4 (1): 27-43.
- [3] Bernhardt OH, S.; Sümnick, W.; Meyer, G. (2012) Electrical stimulation of the temporalis muscle during sleep of myofascial pain - a pilot study. *J CranioMand Func* 4 (3): 191–193.
- [4] Boselli M, Parrino L, Smerieri A, Terzano MG (1998) Effect of age on EEG arousals in normal sleep. *Sleep* 21 (4):351–357.
- [5] Clark GT, Beemsterboer PL, Solberg WK, Rugh JD (1979) Nocturnal electromyographic evaluation of myofascial pain dysfunction in patients undergoing occlusal splint therapy. *J Am Dent Assoc* 99 (4):607–611.
- [6] Clark GT, Seligman DA, Solberg WK, Pullinger AG (1989) Guidelines for the examination and diagnosis of temporomandibular disorders. *J Craniomandib Disord* 3 (1):7–14.
- [7] EEG arousals: scoring rules and examples: a preliminary report from the Sleep Disorders Atlas Task Force of the American Sleep Disorders Association (1992). *Sleep* 15 (2):173–184.
- [8] Feu D, Catharino F, Quintao CCA, De Oliveira Almeida MA (2013) A systematic review of etiological risk factors associated with bruxism. *Journal of Orthodontics* 40 (2):163–171.
- [9] Glaros AG (1981) Incidence of diurnal and nocturnal bruxism. *J Prosthet Dent* 45 (5):545–549.
- [10] Harada T, Ichiki R, Tsukiyama Y, Koyano K (2006) The effect of oral splint devices on sleep bruxism: a 6-week observation with an ambulatory electromyographic recording device. *J Oral Rehabil* 33 (7):482–488.
- [11] Huynh N, Lavigne GJ, Lanfranchi PA, Montplaisir JY, de Champlain J (2006) The effect of 2 sympatholytic medications--propranolol and clonidine-on sleep bruxism: experimental randomized controlled studies. *Sleep* 29 (3):307–316.
- [12] Huynh NT, Rompre PH, Montplaisir JY, Manzini C, Okura K, Lavigne GJ (2006) Comparison of various treatments for sleep bruxism using determinants of number needed to treat and effect size. *Int J Prosthodont* 19 (5):435–441.
- [13] Jadidi F, Castrillon E, Svensson P (2008) Effect of conditioning electrical stimuli on temporalis electromyographic activity during sleep. *J Oral Rehabil* 35 (3):171–183.
- [14] Jadidi F, Castrillon EE, Nielsen P, Baad-Hansen L, Svensson P (2013) Effect of contingent electrical stimulation on jaw muscle activity during sleep: a pilot study with a randomized controlled trial design. *Acta Odontol Scand* 71 (5):1050–1062.
- [15] Karadeniz D, Ondze B, Besset A, Billiard M (2000) EEG arousals and awakenings in relation with periodic leg movements during sleep. *J Sleep Res* 9 (3):273–277.
- [16] Khoury S, Rouleau GA, Rompre PH, Mayer P, Montplaisir JY, Lavigne GJ (2008) A significant increase in breathing amplitude precedes sleep bruxism. *Chest* 134 (2):332–337.
- [17] Landry-Schonbeck A, de Grandmont P, Rompre PH, Lavigne GJ (2009) Effect of an adjustable mandibular advancement appliance on sleep bruxism: a crossover sleep laboratory study. *Int J Prosthodont* 22 (3):251–259.
- [18] Lavigne GJ, Khoury S, Abe S, Yamaguchi T, Raphael K (2008) Bruxism physiology and pathology: an overview for clinicians. *J Oral Rehabil* 35 (7):476–494.

- [19] Lavigne GJ, Soucy JP, Lobbezoo F, Manzini C, Blanchet PJ, Montplaisir JY (2001) Double-blind, crossover, placebo-controlled trial of bromocriptine in patients with sleep bruxism. *Clin Neuropharmacol* 24 (3):145–149.
- [20] Lobbezoo F, Ahlberg J, Glaros AG, Kato T, Koyano K, Lavigne GJ, de Leeuw R, Manfredini D, Svensson P, Winocur E (2013) Bruxism defined and graded: an international consensus. *J Oral Rehabil* 40 (1):2–4.
- [21] Lobbezoo F, Ahlberg J, Manfredini D, Winocur E (2012) Are bruxism and the bite causally related? *J Oral Rehabil* 39 (7):489–501.
- [22] Lobbezoo F, Lavigne GJ, Tanguay R, Montplaisir JY (1997) The effect of catecholamine precursor L-dopa on sleep bruxism: a controlled clinical trial. *Mov Disord* 12 (1):73–78.
- [23] Lobbezoo F, Rompre PH, Soucy JP, Iafrancesco C, Turkewicz J, Montplaisir JY, Lavigne GJ (2001) Lack of associations between occlusal and cephalometric measures, side imbalance in striatal D2 receptor binding, and sleep-related oromotor activities. *J Orofac Pain* 15 (1):64–71.
- [24] Macaluso GM, Guerra P, Di Giovanni G, Boselli M, Parrino L, Terzano MG (1998) Sleep bruxism is a disorder related to periodic arousals during sleep. *J Dent Res* 77 (4):565–573.
- [25] Madani AS, Abdollahian E, Khiavi HA, Radvar M, Foroughipour M, Asadpour H, Hasanzadeh N (2013) The efficacy of gabapentin versus stabilization splint in management of sleep bruxism. *J Prosthodont* 22 (2):126–131.
- [26] Magee KR (1970) Bruxism related to levodopa therapy. *Jama* 214 (1):147.
- [27] Maluly M, Andersen ML, Dal-Fabbro C, Garbuio S, Bittencourt L, de Siqueira JT, Tufik S (2013) Polysomnographic study of the prevalence of sleep bruxism in a population sample. *J Dent Res* 92 (7 Suppl):97s–103s.
- [28] Manfredini D, Landi N, Fantoni F, Segu M, Bosco M (2005) Anxiety symptoms in clinically diagnosed bruxers. *J Oral Rehabil* 32 (8):584–588.
- [29] Manfredini D, Winocur E, Guarda-Nardini L, Paesani D, Lobbezoo F (2013) Epidemiology of bruxism in adults: a systematic review of the literature. *J Orofac Pain* 27 (2):99–110.
- [30] Mohamed SE, Christensen LV, Penchas J (1997) A randomized double-blind clinical trial of the effect of amitriptyline on nocturnal masseteric motor activity (sleep bruxism). *Cranio* 15 (4):326–332.
- [31] Nishi SE, Basri R, Alam MK (2016) Uses of electromyography in dentistry: An overview with meta-analysis. *Eur J Dent* 10 (3):419–425.
- [32] Pierce CJ, Chrisman K, Bennett ME, Close JM (1995) Stress, anticipatory stress, and psychologic measures related to sleep bruxism. *J Orofac Pain* 9 (1):51–56.
- [33] Ramar K, Dort LC, Katz SG, Lettieri CJ, Harrod CG, Thomas SM, Chervin RD (2015) Clinical Practice Guideline for the Treatment of Obstructive Sleep Apnea and Snoring with Oral Appliance Therapy: An Update for 2015. *J Clin Sleep Med* 11 (7): 773–827.
- [34] Rehm DD, Mainieri VC, Saueressig AC, Grossi PK, Teixeira ER, Tenenbaum HC, Drummond LG, Grossi ML (2012) Effects of the bite splint 15-day treatment termination in patients with temporomandibular disorder with a clinical history of sleep bruxism: a longitudinal single-cohort study. *Oral Surg Oral Med Oral Pathol Oral Radiol* 114 (6): 740–748.
- [35] Rugh JD, Solberg WK (1975) Electromyographic studies of bruxist behavior before and during treatment. *J Calif Dent Assoc* 3 (9): 56–59.
- [36] van der Zaag J, Lobbezoo F, Wicks DJ, Visscher CM, Hamburger HL, Naeije M (2005) Controlled assessment of the efficacy of occlusal stabilization splints on sleep bruxism. *J Orofac Pain* 19 (2): 151–158.
- [37] Watanabe T, Ichikawa K, Clark GT (2003) Bruxism levels and daily behaviors: 3 weeks of measurement and correlation. *J Orofac Pain* 17 (1): 65–73.
- [38] Winocur E, Gavish A, Voikovitch M, Emodi-Perlman A, Eli I (2003) Drugs and bruxism: a critical review. *J Orofac Pain* 17 (2): 99–111.

- [1] Tahmaseb A, Wismeijer D, Coucke W, Derksen W. Computer technology applications in surgical implant dentistry: a systematic review. *Int J Oral Maxillofac Implants*. 2014.
- [2] Van Assche N, Vercruyssen M, Coucke W, Teughels W, Jacobs R, Quirynen M. Accuracy of computer-aided implant placement. *Clin Oral Implants Res*. 2012 Oct.
- [3] Vercruyssen M, Cox C, Coucke W, Naert I, Jacobs R, Quirynen M. A randomized clinical trial comparing guided implant surgery (bone- or mucosa-supported) with mental navigation or the use of a pilot-drill template. *J Clin Periodontol*. 2014 Jul.
- [4] Gan N, Xiong Y, Jiao T. Accuracy of Intraoral Digital Impressions for Whole Upper Jaws, Including Full Dentitions and Palatal Soft Tissues. *PLoS One*. 2016 Jul.
- [5] Aragón ML, Pontes LF, Bichara LM, Flores-Mir C, Normando D. Validity and reliability on introral scanners compared to conventional gypsum models measurements: a systematic review. *Eur J Orthod*. 2016 Aug.
- [6] Ekfeldt A, Furst B, Carlsson GE. Zirconia abutments for single-tooth implant restorations: a 10- to 11-year follow-up study. *Clin Oral Implants Res*. 2016 Sep.
- [7] De Avila ED, Avila-Campos MJ, Vergani CE, Spolidório DM, Mollo Fde A Jr. Structural and quantitative analysis of a mature anaerobic biofilm on different implant abutment surfaces. *J Prosthet Dent*. 2016 Apr.
- [8] Spies BC, Witkowski S, Butz F, Vach K, Kohal RJ. Bi-layered zirconia/fluor-apatite bridges supported by ceramic dental implants: a prospective case series after thirty month of observation. *Clin Oral Implants Res*. 2016 Oct.
- [9] Ioannidis A, Bindl A. Clinical prospective evaluation of zirconia-based three-unit posterior fixed dental protheses: Up-to ten-year results. *J Dent*. 2016 Apr.
- [10] Park JH, Park S, Lee K, Yun KD, Lim HP. Antagonist wear of three CAD/CAM anatomic contour zirconia ceramics. *J Prosthet Dent*. 2014 Jan.
- [11] Miyazaki T, Nakamura T, Matsumura H, Ban S, Kobayashi T. Current status of zirconia restorations. *J Prosthodont Res*. 2013 Oct.
- [12] Staubli N, Walter C, Schmidt JC, Weiger R, Zitzmann NU. Excess cement and the risk of peri-implant disease – a systematic review. *Clin Oral Implants Res*.