

## **ZMK 34 (1–2) 2018**

**Prof. Dr. Jürgen Manhart**

### **Seitenzahnrestauration mitOrmoceren – ein klinischer Fallbericht**

- [1] Kunzelmann KH: Komposite – komplexe Wunder moderner Dentaltechnologie. Teil 1: Füllkörpertechnologie. *Ästhetische Zahnmedizin* 10 (3), 14–24 (2007).
- [2] Kunzelmann KH: Komposite – komplexe Wunder moderner Dentaltechnologie. Teil 2: Matrixchemie. *Ästhetische Zahnmedizin* 11 (1), 22–35 (2008).
- [3] Ferracane JL: Resin composite – state of the art. *Dent Mater* 27 (1), 29–38 (2011).
- [4] Manhart J: Charakterisierung direkter zahnärztlicher Füllungsmaterialien für den Seitenzahnbereich. Alternativen zum Amalgam? *Quintessenz* 57 (5), 465–481 (2006).
- [5] Vanini L: Light and color in anterior composite restorations. *Pract Periodontics Aesthet Dent* 8 (7), 673–682 (1996).
- [6] Ardu S, et al.: Influence of water sorption on resin composite color and color variation amongst various composite brands with identical shade code: an in vitro evaluation. *J Dent* 39 Suppl 1, e37–44 (2011).
- [7] Manhart J: Direkte Restaurationen im Frontzahnbereich (Teil 1) – Möglichkeiten moderner Kompositsysteme. *Zahnärzteblatt Baden-Württemberg* 44 (7), 8–14 (2016).
- [8] Manhart J, Hickel R: „Bulk Fill“-Komposite. Neuartige Einsatztechnik von Kompositen im Seitenzahnbereich. *Swiss Dental Journal* 124 (1), 19–28 (2014).
- [9] Manhart J: Bulk-fill composites for posterior restorations. *Inside Dentistry* 11 (3), 58–65 (2015).
- [10] Lenhard M: Bulk-Fill-Komposite – der schnelle Weg zur Restauration? *Quintessenz* 66 (8), 879–895 (2015).
- [11] Manhart J: Muss es immer Kaviar sein? – Die Frage nach dem Aufwand für Komposite im Seitenzahnbereich. *ZMK 27* (Sonderausgabe März 2011), 10–15 (2011).
- [12] Christensen GJ: Advantages and challenges of bulk-fill resins. *Clinicians Report* 5 (1), 1–2 (2012).
- [13] Noack MJ: Wirtschaftliche Füllungstechnik. Warum und womit? *ZWR Das Deutsche Zahnärzteblatt* 122 (3), 86–94 (2013).
- [14] Margeas RC: Bulk-fill materials: simplify restorations, reduce chairtime. *cCompend Contin Educ Dent* 36 (1), e1–e4 (2015).
- [15] Ilie N, Stawarczyk B: Bulk-Fill-Komposite: neue Entwicklungen oder doch herkömmliche Komposite? *ZMK 30* (3), 90–97 (2014).
- [16] Tauböck TT: Bulk-Fill-Komposite. Wird die Füllungstherapie einfacher, schneller und erfolgreicher? *teamwork J Cont Dent Educ* 16 (4), 318–323 (2013).
- [17] Ferracane JL, Greener EH: The effect of resin formulation on the degree of conversion and mechanical properties of dental restorative resins. *J Biomed Mater Res* 20 (1), 121–131 (1986).
- [18] Caughman WF, et al.: Correlation of cytotoxicity, filler loading and curing time of dental composites. *Biomaterials* 12 (8), 737–740 (1991).
- [19] Feilzer AJ, de Gee AJ, Davidson CL: Setting stress in composite resin in relation to configuration of the restoration. *J Dent Res* 66, 1636–1639 (1987).
- [20] Federlin M, et al.: Kompositrestaurationen im Seitenzahnbereich. S1-Handlungsempfehlung (Langversion). AWMF-Registernummer: 083–028; Stand: Oktober 2016; gültig bis: Oktober 2021. *Deutsche Zahnärztliche Zeitschrift* 72 (1), 75–82 (2017).

- [21] Margeas R: New bulk-fill material simplifies restorations to one step. *Inside Dentistry* 10 (10), 86–90 (2014).
- [22] Burtscher P: Von geschichteten Inkrementen zur Vier-Millimeter-Bulk-Fill-Technik – Anforderungen an Komposit und Lichthärtung. *DZW Die Zahnarzt Woche* 39, 6–8 (2011).
- [23] Czasch P, Ilie N: In vitro comparison of mechanical properties and degree of cure of bulk fill composites. *Clin Oral Investig* 17 (1), 227–235 (2013).
- [24] Finan L, et al.: The influence of irradiation potential on the degree of conversion and mechanical properties of two bulk-fill flowable RBC base materials. *Dent Mater* 29 (8), 906–12 (2013).
- [25] Manhart J: Neues Konzept zum Ersatz von Dentin in der kompositbasierten Seitenzahnversorgung. *ZWR Das Deutsche Zahnärzteblatt* 119 (3), 118–125 (2010).
- [26] Fleming GJ, et al.: The potential of a resin-composite to be cured to a 4mm depth. *Dental Materials* 24 (4), 522–529 (2008).
- [27] Ilie N, Kessler A, Durner J: Influence of various irradiation processes on the mechanical properties and polymerisation kinetics of bulk-fill resin based composites. *J Dent* 41 (8), 695–702 (2013).
- [28] Ferracane J, Alex G, Margeas R: Question: Are bulk-fill composites a good idea? *Inside Dentistry* 10 (10), 42–44 (2014).
- [29] Hickel R: Neueste Komposite – viele Behauptungen. *BZB Bayerisches Zahnärzteblatt* 49 (9), 50–53 (2012).
- [30] Ilie N, Bucuta S, Draenert M: Bulk-fill resin-based composites: an in vitro assessment of their mechanical performance. *Oper Dent* 38 (6), 618–625 (2013).
- [31] Condon JR, Ferracane JL: Evaluation of composite wear with a new multi-mode oral wear simulator. *Dent Mater* 12 (4), 218–226 (1996).
- [32] Condon JR, Ferracane JL: In vitro wear of composite with varied cure, filler level, and filler treatment. *J Dent Res* 76 (7), 1405–1411 (1997).
- [33] Poggio C, et al.: Surface roughness of flowable resin composites eroded by acidic and alcoholic drinks. *J Conserv Dent* 15 (2), 137–140 (2012).
- [34] Frankenberger R, Garcia-Godoy F, Kramer N: Clinical performance of viscous glass ionomer cement in posterior cavities over two years. *Int J Dent*, p. Article-ID: 781462 (2009).
- [35] Lohbauer U: Dental glass ionomer cements as permanent filling materials? – Properties, limitations and future trends. *Materials* 3 (1), 76–96 (2010).
- [36] Hickel R, et al.: Direkte Kompositrestaurationen im Seitenzahnbereich – Indikation und Lebensdauer. Gemeinsame Stellungnahme der Deutschen Gesellschaft für Zahnerhaltung (DGZ) und der Deutschen Gesellschaft für Zahn-, Mund- und Kieferheilkunde (DGZMK) aus dem Jahr 2005. *Deutsche Zahnärztliche Zeitschrift* 60 (10), 543–545 (2005).
- [37] Burke FJ, Lucarotti PS: Re-intervention in glass ionomer restorations: what comes next? *J Dent* 37 (1), 39–43 (2009).
- [38] Scholtanus JD, Huysmans MC: Clinical failure of class-II restorations of a highly viscous glass-ionomer material over a 6-year period: a retrospective study. *J Dent* 35 (2), 156–162 (2007).
- [39] Klink T, et al.: Clinical performance during 48 months of two current glass ionomer restorative systems with coatings: a randomized clinical trial in the field. *Trials* 17 (1), 239 (2016).
- [40] Frankenberger R, et al.: Bulk-Fill-Komposite: Mit dicken Schichten einfacher zum Erfolg? *Quintessenz* 65 (5), 579–584 (2012).
- [41] Frankenberger R, et al.: Die richtige Basisversorgung – Expertenzirkel. *Dental Magazin* 30 (1), 12–24 (2012).

- [42] Peutzfeldt A: Resin composites in dentistry: the monomer systems. *Eur J Oral Sci* 105 (2), 97–116 (1997).
- [43] Guggenberger R, Weinmann W: Exploring beyond methacrylates. *American Journal of Dentistry* 13 (Special Issue), 82–84 (2000).
- [44] Weinmann W, Thalacker C, Guggenberger R: Siloranes in dental composites. *Dent Mater* 21 (1), 68–74 (2005).
- [45] Lien W, Vandewalle KS: Physical properties of a new silorane-based restorative system. *Dent Mater* 26 (4), 337–344 (2010).
- [46] Ilie N, Hickel R: Silorane-based dental composite: behavior and abilities. *Dent Mater J* 25 (3), 445–454 (2006).
- [47] Ilie N, Hickel R: Macro-, micro- and nano-mechanical investigations on silorane and methacrylate-based composites. *Dent Mater* 25 (6), 810–819 (2009).
- [48] Zimmerli B, et al.: Composite materials: composition, properties and clinical applications. A literature review. *Schweiz Monatsschr Zahnmed* 120 (11), 972–986 (2010).
- [49] Manhart J, et al.: Randqualität von Ormocer- und Kompositfüllungen in Klasse-II-Kavitäten nach künstlicher Alterung. *Deutsche Zahnärztliche Zeitschrift* 54, 89–95 (1999).
- [50] Wolter H, Storch W: Neuartige Silanklasse – Werkstoffe für Formkörper. *ISC-Tätigkeitsbericht* 1992, 61–72 (1992).
- [51] Wolter H, Storch W, Ott H: Dental filling materials (posterior composites) based on inorganic/organic copolymers (ORMOCERs). *MACRO AKRON* (35th IUPAC, International Symposium on Macromolecules, Ohio, Proceedings), 503 (1994).
- [52] Wolter H, Storch W, Ott H: New inorganic/organic copolymers (ORMOCERs) for dental applications. *Materials Research Society Symposia Proceedings* 346, 143–149 (1994).
- [53] Wolter H: Kompakte Ormocere und Ormocer-Komposite. *Fraunhofer-Institut für Silikatforschung (ISC) – Tätigkeitsbericht* 1995, 56–63 (1995).
- [54] Wolter H, et al.: Neue biokompatible Dentalwerkstoffe auf Ormocer-Basis. In: Planck H, Stallforth H (Eds.); *Tagungsband Werkstoffwoche 1998, Band 4, Symposium 4: Werkstoffe für die Medizintechnik*. Wiley VCH: Weinheim. 245–248 (1998).
- [55] Manhart J, et al.: Mechanical properties and wear behavior of light-cured packable composite resins. *Dental Materials* 16, 33–40 (2000).
- [56] Hickel R, et al.: New direct restorative materials. *FDI Commission Project. Int Dent J* 48 (1), 3–16 (1998).
- [57] Greiwe K, Schottner G: ORMOCERe: Eine neue Werkstoffklasse. *FhG-Berichte* 2, 64–67 (1990).
- [58] Moszner N, et al.: Sol-Gel Materials, 1. Synthesis and hydrolytic condensation of new cross-linking alkoxysilane methacrylates and light-curing composites based upon the condensates. *Macromol Mater Eng* 287 (5), 339–347 (2002).
- [59] Moszner N, et al.: Sol-gel materials 2. Light-curing dental composites based on ormocers of cross-linking alkoxysilane methacrylates and further nano-components. *Dent Mater* 24 (6), 851–856 (2008).
- [60] Wolter H: Werkstoffe mit Biss. Teil I: Ormocere. *DZW Die Zahnarzt Woche* 11/15, 10–11 (2015).
- [61] Ilie N, Hickel R: Resin composite restorative materials. *Aust Dent J* 56 (Suppl 1), 59–66 (2011).
- [62] de Goes MF, Shinohara MS, Freitas MS: Performance of a new one-step multi-mode adhesive on etched vs non-etched enamel on bond strength and interfacial morphology. *J Adhes Dent* 16 (3), 243–250 (2014).
- [63] Hanabusa M, et al.: Bonding effectiveness of a new 'multi-mode' adhesive to enamel and dentine. *J Dent* 40 (6), 475–84 (2012).

- [64] McLean DE, et al.: Enamel bond strength of new universal adhesive bonding agents. *Oper Dent* 40 (4), 410–417 (2015).
- [65] Takamizawa T, et al.: Influence of different etching modes on bond strength and fatigue strength to dentin using universal adhesive systems. *Dent Mater* 32 (2), e9–21 (2016).
- [66] Wagner A, et al.: Bonding performance of universal adhesives in different etching modes. *J Dent* 42 (7), 800–807 (2014).
- [67] Lenzi TL, et al.: Bonding performance of a multimode adhesive to artificially-induced caries-affected primary dentin. *J Adhes Dent* 17 (2), 125–131 (2015).
- [68] Loguercio AD, et al.: A new universal simplified adhesive: 36-Month randomized double-blind clinical trial. *J Dent* 43 (9), 1083–1092 (2015).
- [69] Munoz MA, et al.: In vitro longevity of bonding properties of universal adhesives to dentin. *Oper Dent* 40 (3), 282–292 (2015).
- [70] Da Rosa Rodolpho PA, et al.: 22-year clinical evaluation of the performance of two posterior composites with different filler characteristics. *Dent Mater* 27 (10), 955–963 (2011).
- [71] van de Sande FH, et al.: 18-year survival of posterior composite resin restorations with and without glass ionomer cement as base. *Dent Mater* 31 (6), 669–675 (2015).
- [72] Manhart J, et al.: Review of the clinical survival of direct and indirect restorations in posterior teeth of the permanent dentition. *Oper Dent* 29 (5), 481–508 (2004).
- [73] Heintze SD, Rousson V: Clinical effectiveness of direct class II restorations – a meta-analysis. *J Adhes Dent* 14 (5), 407–431 (2012).
- [74] Opdam NJ, et al.: Longevity of posterior composite restorations: a systematic review and meta-analysis. *J Dent Res* 93 (10), 943–949 (2014).
- [75] Opdam NJ, et al.: 12-year survival of composite vs. amalgam restorations. *J Dent Res* 89 (10), 1063–1067 (2010).
- [76] Pallesen U, van Dijken JW: A randomized controlled 30 years follow up of three conventional resin composites in Class II restorations. *Dent Mater* 31 (10), 1232–1244 (2015).
- [77] Pallesen U, van Dijken JW: A randomized controlled 27 years follow up of three resin composites in Class II restorations. *J Dent* 43 (12), 1547–1558 (2015).

- [1] Ahovuo-Saloranta A, Forss H, Walsh T, Nordblad A, Makela M, Worthington HV: Pit and fissure sealants for preventing dental decay in permanent teeth. The Cochrane database of systematic reviews 7, CD001830 (2017). doi:10.1002/14651858.CD001830.pub5
- [2] Weintraub JA, Burt BA: Prevention of dental caries by the use of pit-and-fissure sealants. Journal of public health policy 8 (4), 542-560 (1987).
- [3] Liebenberg WH: The fissure sealant impasse. Quintessence international 25 (11), 741-745 (1994).
- [4] Muller-Bolla M, Lupi-Pegurier L, Tardieu C, Velly AM, Antomarchi C: Retention of resin-based pit and fissure sealants: A systematic review. Community dentistry and oral epidemiology 34 (5), 321-336 (2006). doi:10.1111/j.1600-0528.2006.00319.x
- [5] Kuhnisch J, Mansmann U, Heinrich-Weltzien R, Hickel R: Longevity of materials for pit and fissure sealing--results from a meta-analysis. Dental materials : official publication of the Academy of Dental Materials 28 (3), 298-303 (2012). doi:10.1016/j.dental.2011.11.002
- [6] Mickenautsch S, Yengopal V: Validity of sealant retention as surrogate for caries prevention -- a systematic review. PloS one 8 (10), e77103 (2013). doi:10.1371/journal.pone.0077103
- [7] Mickenautsch S, Yengopal V: Retention loss of resin based fissure sealants - a valid predictor for clinical outcome? The open dentistry journal 7, 102-108 (2013). doi:10.2174/18742106201305130001
- [8] Wright JT, Tampi MP, Graham L, Estrich C, Crall JJ, Fontana M, Gillette EJ, Novy BB, Dhar V, Donly K, Hewlett ER, Quinonez RB, Chaffin J, Crespin M, Iafolla T, Siegal MD, Carrasco-Labra A: Sealants for preventing and arresting pit-and-fissure occlusal caries in primary and permanent molars: A systematic review of randomized controlled trials-a report of the American Dental Association and the American Academy of Pediatric Dentistry. Journal of the American Dental Association 147 (8), 631-645 e618 (2016). doi:10.1016/j.adaj.2016.06.003
- [9] Schwendicke F, Jager AM, Paris S, Hsu LY, Tu YK: Treating pit-and-fissure caries: a systematic review and network meta-analysis. Journal of dental research 94 (4), 522-533 (2015). doi:10.1177/0022034515571184
- [10] 2016 SLF S3 Leitlinie zur Fissuren- und Grübchenversiegelung, Langversion 2016. AWMF Registernummer: 083/002. <http://www.awmf.org/leitlinien/detail/II/083-002.html>. Accessed 20.07.17 2017
- [11] Ehrental JC, Graetz C, Plaumann A, Dorfer CE, Herzog W: Number of teeth predict depressive symptoms in a longitudinal study on patients with periodontal disease. Journal of psychosomatic research 89, 16-19 (2016). doi:10.1016/j.jpsychores.2016.08.002
- [12] Kolavic Gray S, Griffin SO, Malvitz DM, Gooch BF: A comparison of the effects of toothbrushing and handpiece prophylaxis on retention of sealants. Journal of the American Dental Association (1939) 140 (1), 38-46 (2009).
- [13] Griffin SO, Jones K, Gray SK, Malvitz DM, Gooch BF: Exploring four-handed delivery and retention of resin-based sealants. Journal of the American Dental Association (1939) 139 (3), 281-289 (2008); quiz 358

- [14] Bagherian A, Sarraf Shirazi A: Preparation before acid etching in fissure sealant therapy: yes or no?: A systematic review and meta-analysis. *Journal of the American Dental Association* 147 (12), 943-951 (2016). doi:10.1016/j.adaj.2016.08.003
- [15] Bagherian A, Sarraf Shirazi A, Sadeghi R: Adhesive systems under fissure sealants: yes or no?: A systematic review and meta-analysis. *Journal of the American Dental Association* (1939) 147 (6), 446-456 (2016). doi:10.1016/j.adaj.2016.01.014