

Wirksamkeit noninvasiver Therapien zur Inaktivierung oder Reduktion der Wurzelkaries

- [1] Baca P, et al: Effect of chlorhexidine-thymol varnish on root caries in a geriatric population: a randomized double-blind clinical trial. *J Dent* 37 (9), 679–85 (2009).
- [2] Banoczy J, Nemes J: Effect of amine fluoride (AmF)/stannous fluoride (SnF₂) toothpaste and mouthwashes on dental plaque accumulation, gingivitis and root-surface caries. *Proc Finn Dent Soc* 87 (4), 555–9 (1991).
- [3] Banting DW, et al: The effectiveness of 10 % chlorhexidine varnish treatment on dental caries incidence in adults with dry mouth. *Gerodontology* 17 (2), 67–76 (2000).
- [4] Baysan A, Lynch E: Clinical reversal of root caries using ozone: 6-month results. *Am J Dent* 20 (4), 203–8 (2007).
- [5] Baysan A, et al: Reversal of primary root caries using dentifrices containing 5,000 and 1,100 ppm fluoride. *Caries Res* 35 (1), 41–6 (2001).
- [6] Davey Smith G, Egger M: Meta-analysis. Unresolved issues and future developments. *BMJ* 316 (7126), 221–5 (1998).
- [7] DePaola PF: Caries in our aging population: What are we learning?, in *Cariology for the nineties*, Bowden GH and Tabak LA, Editors. University of Rochester Press: Rochester, NY. 25–35 (1993).
- [8] Ekstrand KR, et al: A Randomized Clinical Trial of the Anti-Caries Efficacy of 5,000 Compared to 1,450 ppm Fluoridated Toothpaste on Root Caries Lesions in Elderly Disabled Nursing Home Residents. *Caries Res* 47 (5), 391–398 (2013).
- [9] Ettinger RL: Epidemiology of dental caries. A broad review. *Dent Clin North Am* 43 (4), 679–94 (1999).
- [10] Fure S, Gahnberg L, Birkhed D: A comparison of four home-care fluoride programs on the caries incidence in the elderly. *Gerodontology* 15 (2), 51–60 (1998).
- [11] Fure S, Lingstrom P: Evaluation of different fluoride treatments of initial root carious lesions in vivo. *Oral Health Prev Dent* 7 (2), 147–54 (2009).
- [12] Geiger AM, et al: Reducing white spot lesions in orthodontic populations with fluoride rinsing. *Am J Orthod Dentofacial Orthop* 101 (5), 403–7 (1992).
- [13] Griffin SO, et al: Estimating Rates of New Root Caries in Older Adults. *J Dent Res* 83 (8), 634–638 (2004).
- [14] Guyatt GH, et al: GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ* 336 (7650), 924–6 (2008).
- [15] Higgins JPT, Green SG: *Cochrane Handbook for Systematic Reviews of Interventions*. Version 5. 10 [updated March 2011]. The Cochrane Collaboration. Available from www.cochrane-handbook.org. 2011.
- [16] Holmes J: Clinical reversal of root caries using ozone, double-blind, randomised, controlled 18-month trial. *Gerodontology* 20 (2), 106–14 (2003).
- [17] Hu DY, et al: A clinical investigation of the efficacy of a dentifrice containing 1.5 % arginine and 1450 ppm fluoride, as sodium monofluorophosphate in a calcium base, on primary root caries. *J Clin Dent* 24 Spec no A, A23–31 (2013).
- [18] Jensen ME, Kohout F: The effect of a fluoridated dentifrice on root and coronal caries in an older adult population. *J Am Dent Assoc* 117 (7), 829–32 (1988).

- [19] Marinho VC, et al: Topical fluoride (toothpastes, mouthrinses, gels or varnishes) for preventing dental caries in children and adolescents. *Cochrane Database Syst Rev* (4), CD002782 (2003).
- [20] Michaelis W, Schiffner U: The Fourth German Oral Health Study (DMS IV). Institute of German Dentists (IDZ), Deutscher Zahnärzte Verlag: Köln (2006).
- [21] Nitschke I: Zahnmedizinische Grundlagen zur geriatrischen Rehabilitation – eine Einführung in die Alternszahnmedizin. *Zeitschrift für Gerontologie und Geriatrie* 33 (1 Supplement), 45–049 (2000).
- [22] Papas A, et al: Comparative efficacy of stabilized stannous fluoride/sodium hexametaphosphate dentifrice and sodium fluoride/triclosan/copolymer dentifrice for the prevention of periodontitis in xerostomic patients: a 2-year randomized clinical trial. *J Periodontol* 78 (8), 1505–14 (2007).
- [23] Papas A, et al: Caries clinical trial of a remineralising toothpaste in radiation patients. *Gerodontology* 25 (2), 76–88 (2008).
- [24] Paraskevas S, et al: Amine fluoride/stannous fluoride and incidence of root caries in periodontal maintenance patients. A 2 year evaluation. *J Clin Periodontol* 31 (11), 965–71 (2004).
- [25] Peters MC,: Strategies for noninvasive demineralized tissue repair. *Dent Clin North Am* 54 (3), 507–25 (2010).
- [26] Petersson LG, et al: Remineralization of primary root caries lesions using an amine fluoride rinse and dentifrice twice a day. *Am J Dent* 20 (2), 93-6 (2007).
- [27] Powell LV, et al: Caries prevention in a community-dwelling older population. *Caries Res* 33 (5), 333–9 (1999).
- [28] Ravald N, Birkhed D: Prediction of root caries in periodontally treated patients maintained with different fluoride programmes. *Caries Res* 26 (6), 450–8 (1992).
- [29] Ripa LW, et al: Effect of a 0.05 % neutral NaF mouthrinse on coronal and root caries of adults. *Gerodontology* 6 (4), 131–6 (1987).
- [30] Schaeken MJ, Keltjens HM, Van Der Hoeven JS: Effects of fluoride and chlorhexidine on the microflora of dental root surfaces and progression of root-surface caries. *J Dent Res* 70 (2), 150–3 (1991).
- [31] Shellis RP: A scanning electron-microscopic study of solubility variations in human enamel and dentine. *Arch Oral Biol* 41 (5), 473–84 (1996).
- [32] Souza ML, et al: Comparing the efficacy of a dentifrice containing 1.5 % arginine and 1450 ppm fluoride to a dentifrice containing 1450 ppm fluoride alone in the management of primary root caries. *J Dent* 41 Suppl 2, S35–41 (2013).
- [33] Srinivasan M, et al: High-fluoride toothpaste: a multicenter randomized controlled trial in adults. *Community Dent Oral Epidemiol* (2013).
- [34] Tan HP, et al: A randomized trial on root caries prevention in elders. *J Dent Res* 89 (10), 1086–90 (2010).
- [35] US Department of Health and Human Services, Oral Health in America: A Report of the Surgeon General. Rockville, MD: US Department of Health and Human Services, National Institute of Dental and Craniofacial Research, National Institutes of Health, 308 (2000).
- [36] Vered Y, et al: Comparison of a dentifrice containing 0.243 % sodium fluoride, 0.3 % triclosan, and 2.0 % copolymer in a silica base, and a dentifrice containing 0.243 % sodium fluoride in a silica base: a three-year clinical trial of root caries and dental crowns among adults. *J Clin Dent* 20 (2). 62–5 (2009).

- [37] Wallace MC, Retief DH, Bradley EL: The 48-month increment of root caries in an urban population of older adults participating in a preventive dental program. *J Public Health Dent* 53 (3), 133–7 (1993).
- [38] Wierichs RJ, Meyer-Lueckel H: Systematic review on noninvasive treatment of root caries lesions. *J Dent Res* 94 (2), 261–71 (2015).
- [39] Wyatt CC, MacEntee MI: Caries management for institutionalized elders using fluoride and chlorhexidine mouthrinses. *Community Dent Oral Epidemiol* 32 (5), 322–8 (2004).
- [40] Wyatt CC, et al: Chlorhexidine and preservation of sound tooth structure in older adults. A placebo-controlled trial. *Caries Res* 41 (2), 93–101 (2007).
- [41] Zhang W, et al: Silver Diamine Fluoride and Education to Prevent and Arrest Root Caries among Community-Dwelling Elders. *Caries Res* 47 (4), 284–90 (2013).

1: Analyse der Chancen und Grenzen

- [1] Mormann WH: The evolution of the CEREC system. J Am Dent Assoc 137 (Suppl), 7–13 (2006).
- [2] Mormann WH, Brandestini M: Cerec-System: computerized inlays, onlays and shell veneers. Zahnärztl Mitt 77 (21), 2400–2405 (1987).
- [3] Ender A, Attin T, Mehl A; In vivo precision of conventional and digital methods of obtaining complete-arch dental impressions, J Prosthet Dent; 2016; 115(3): 313–320.
- [4] Ender A et al.: In vivo precision of conventional and digital methods for obtaining quadrant dental impressions. Clin Oral Investig 115 (3): 313–320 (2015).
- [5] Ender A, Mehl A: Influence of scanning strategies on the accuracy of digital intraoral scanning systems. Int J Comput Dent 16 (1), 11–21 (2013).
- [6] Ender A, Mehl A: In-vitro evaluation of the accuracy of conventional and digital methods of obtaining full-arch dental impressions. Quintessence Int 46 (1): 9–17. (2014).
- [7] Ender A, Mehl A: Full arch scans: conventional versus digital impressions – an in-vitro study. Int J Comput Dent 14 (1), 11–21 (2011).
- [8] Seelbach P, Brueckel C, Wostmann B: Accuracy of digital and conventional impression techniques and workflow. Clin Oral Investig 17 (7), 1759–1764 (2013).
- [9] Patzelt SB et al.: Accuracy of full-arch scans using intraoral scanners. Clin Oral Investig 18 (6), 1687–1694 (2014).
- [10] Zaruba M, Ender A, Mehl A: New applications for three-dimensional follow-up and quality control using optical impression systems and OraCheck. Int J Comput Dent 17 (1), 53–64 (2014).

Parodontitis – nach der Therapie ist vor der Therapie

- [1] Alaluusua S, Asikainen S, Lai CH: Intrafamilial transmission of *Actinobacillus actinomycetemcomitans*. *Journal of Periodontology* 62 (3), 207–210 (1991).
- [2] Axelsson P, Nyström B, Linde J: The long-term effect of a plaque control program on tooth mortality, caries and periodontal disease in adults. Results after 30 years of maintenance. *J Clin Periodontol* 31 (9), 749–757 (2004).
- [3] Barfuß A: Expertenzirkel. PA-Therapie: Lokale Antibiotika. *Dentalmagazin* 31 (6), 12–28 (2013).
- [4] Eickholz P et al.: Tooth loss after active periodontal therapy. 1: Patient-related factors for risk, prognosis, and quality of outcome. *J Clin Periodontol* 35 (2), 165–174 (2008).
- [5] Eickholz P, Kim TS, Bürklin T, Schacher B, Rengli HH, Schaecken MT, Holle R, Kubler A, Ratka-Krüger P: Non-Surgical periodontal therapy with adjunctive topical doxycycline: a double-blind randomized controlled multicenter study. *J Clin Periodontol* 29 (2), 108–117 (2002).
- [6] Goodson JM: Pharmacokinetic principles controlling efficacy of oral therapy. *J Dent Res* 68 (special issue), 1625–1632 (1989).
- [7] Halling F: Antibiotika in der Zahnmedizin. *Zahnmedizin up2date*, 67-80 (1/2014). Abruf am 29.09.2015:
https://www.thieme.de/statics/dokumente/thieme/final/de/dokumente/tw_zahnmedizin/Antibiotika_Zahnmedizin_up2date_2014_1.pdf
- [8] Interne Daten, Heraeus Kulzer GmbH.
- [9] Jorgensen MG, Slots J: Practical antimicrobial periodontal therapy. *Compend Contin Educ Dent* 21, 111–120 (2000).
- [10] Kim T et al: Pharmacokinetic profile of a locally administered doxycycline gel in crevicular fluid, blood, and saliva. *J Periodontol* 73 (11), 1285–1291 (2002).
- [11] Lang NP, Tonetti MS: Periodontal risk assessment (PRA) for patients in supportive periodontal therapy (SPT). *Oral Health Prevent Dent* 1 (1), 7-16 (2002).
- [12] Lindhe J, Nyman S: Long-term maintenance of patients treated for advanced periodontal disease. *J Clin Periodontol* 11 (8), 504–514 (1984).
- [13] Magnusson I, Lindhe J, Yoneyama T, Liljenberg B: Recolonization of a subgingival microbiota following scaling in deep pockets. *J Clin Periodontol* 11 (3), 193–207 (1984).
- [14] Matesanz-Pérez P, García-Gargallo M, Figuero E, Bascones-Martínez A, Sanz M, Herrera DJ: A systematic review on the effects of local antimicrobials as adjuncts to subgingival debridement, compared with subgingival debridement alone, in the treatment of chronic periodontitis. *Clin Periodontol* 40 (3), 227–241 (2013).
- [15] Nyman S, Lindhe J, Rosling B: Periodontal surgery in plaque-infected dentitions. *J Clin Periodontol* 4 (4), 240–249 (1977).
- [16] Ratka-Krüger P, Schacher B, Bürklin T, Böddinghaus B, Holle R, Renggli HH, Eickholz P, Kim TS: Non-surgical periodontal therapy with adjunctive topical doxycycline: a double-masked, randomized, controlled multicenter study. II. Microbiological results. *J Periodontol* 76 (1), 66–74 (2005).
- [17] Ramfjord SP: Maintenance care for treated periodontitis patients. *J Clin Periodontol* 14 (8), 433–437 (1987).
- [18] Richtlinien des Bundesausschusses der Zahnärzte und Krankenkassen für eine ausreichende, zweckmäßige und wirtschaftliche vertragszahnärztliche Versorgung (Behandlungs-Richtlinien) in der ab 01.01.2004 gültigen Fassung (Stand: 01.01.2005). Darin: B. Vertragszahnärztliche Behandlung. V. Systematische Behandlung von Parodontopathien (Par-Behandlung) Abruf am 29.09.2015:
http://www.kzv-berlin.de/fileadmin/user_upload/Abrechnung/PAR/Richtlinien.pdf

- [19] Slots J, Jorgensen MG: Effective, safe, practical and affordable periodontal antimicrobial therapy: where are we going, and are we there yet? *Periodontol* 28 (1), 298–312 (2000).
- [20] Wissenschaftliche Stellungnahme der DGZMK und DGP: Adjuvante Antibiotika in der Parodontitistherapie. Abruf am 29.09.2015:
http://www.dgzmk.de/uploads/tx_szdgzmkdocuments/Adjuvante_Antibiotika_in_der_Parodontitistherapie.pdf

Moderne CAD/CAM-Kompositmaterialien – deren Materialeigenschaften und Befestigungsstrategien

- [1] Strietzel, R., Lahl, C.: CAD/CAM-Systeme in Labor und Praxis, S. 12 ff. Verlag Neuer Merkur GmbH München (2007), 978-3-937346-41-0 (ISBN).
- [2] Rinke, S.: VOLLKERAMIK – Ein Praxiskonzept, S. 7 ff. Quintessenz Verlag Berlin (2011), 978-3-86867-049-3 (ISBN).
- [3] Gebrauchsanleitung LavaTM Ultimate CAD/CAM Restorationsmaterial, Deutsch, 3M ESPE Dental Products (6/2015), 44-0007-7648-2-A.
- [4] Gebrauchsanleitung CERASMARTTM, Deutsch, GC DENTAL Products Corp (6/2015), 13008790 290655DK.
- [5] Gebrauchsanleitung SHOFU BLOCK HC, Deutsch, SHOFU Inc (1/2014), 09 SS000032.
- [6] Gebrauchsanleitung BRILLIANT Crios (1/2016), 3000399801.16.
- [7] VITA CAD/CAM MATERIALIEN – Alle Lösungen auf einen Blick, Kompendium für Praxen und Labore. 994D – 0115 (X.) S – Version (01), VITA Zahnfabrik H. Rauter GmbH & Co. KG.
- [8] Böhner, R., Claude, M., Kopfmann, C.: Characteristic of polymer based CAD/CAM blocks for permanent restorations, #597, IADR (2015), Antalya, Turkey.
- [9] Spears, I.R., van Noort, R., Crompton, R.H., Cardew, G.E., Howard, I.C.: The Effects of Enamel Anisotropy on the Distribution of Stress in a Tooth. J Dent Res 72, 1526–1531 (1993).
- [10] Meredith, N., Sherriff, M., Setchell, D.J., Swanson, S.A.: Measurement of the microhardness and Young's modulus of human enamel and dentine using an indentation technique, Arch Oral Biol 41, 539–545 (1996).
- [11] Hülsmann, M.: Checklisten der Zahnmedizin, Endodontie, S. 19 ff, Georg Thieme Verlag (2008), 9783131382511 (ISBN).
- [12] Zhi, L., Bortolotto, T., Krejci, I.: Comparative in vitro wear resistance of CAD/CAM composite resin and ceramic materials J Prosthet Dent (2015), pii: S0022-3913(15)00437-0. doi: 10.1016/j.prosdent.2015.07.011. [Epub ahead of print].
- [13] Stawarczyk, B., Liebermann, A., Eichberger, M., Güth, J.-F.: Evaluation of mechanical an optical behaviour of current esthetic dental restorative CAD/CAM composites. J Mech Behav Biomed Mater 55, 1–11 (2015).
- [14] Behr, M., Proff, P., Kolbeck, C., Langrieger, S., Kunze, J., Handel, G., Rosentritt, M.: The bond strength of the resin-to-zirconia interface using different bonding concepts. J Mech Behav Biomed Mater 4, 2–8 (2011).
- [15] Kassotakis, E., Stavridakis, M., Bortolotto, T., Ardu, S., Krejci I.: Evaluation of the Effect of Different Surface Treatments on Luting CAD/CAM Composite Resin Overlay Workpieces. J Adhes Dent 17, 521–528 (2015).
- [16] Frankenberger, R., Hartmann, V.E., Krech, M., Krämer, N., Reich, S., Braun, A., Roggendorf M.: Adhesive luting of new CAD/CAM materials. Int J Comput Dent 18, 9–20 (2015).
- [17] Stawarczyk, B., Stich, N., Eichberger, M., Edelhoff, D., Roos, M., Gernet, W., Keul, C.: Long-term tensile bond strength of differently cemented nanocomposite CAD/CAM crowns on dentin abutment. Dent Mater 30, 334–342 (2014).
- [18] Keul, C., Müller-Hahl, M., Eichberger, M., Liebermann, A., Roos, M., Edelhoff, D., Stawarczyk, B.: Impact of different adhesives on work of adhesion between CAD/CAM polymers and resin composite cements. J Dent 42, 1105–1114 (2014).
- [19] Stawarczyk, B., Krawczuk, A., Ilie, N.: Tensile bond strength of resin composite repair in vitro using different surface preparation conditionings to an aged CAD/CAM resin nanoceramic. Clin Oral Investig 19, 299–308 (2015).

- [20] Zuryati, A.B., Wahyuni, J., Siew, F., Zaihan, A., Dasmawati, M.: Shear bond strength of computer-aided design and computer-aided manufacturing feldspatic and nano resin ceramics blocks cemented with three different generations of resin cement. *Conserv Dent* 18, 355–359 (2015).
- [21] Lührs, A.-K., Pongprueska, P., De Munck, J., Geurtsen, W., Van Meerbeek, B.: Curing mode affects bond strength of adhesively luted composite CAD/CAM restorations to dentin. *Dental Materials* 30, 281–291 (2014).
- [22] Spitznagel, F.A., Horvath, S.D., Guess, P.C., Blatz, M.B.: Resin Bond to Indirect Composite and New Ceramic/Polymer Materials: A Review of the Literature. *J Esth Rest Dent* 26, 382–392 (2014).

40 Jahre Ultracain®: der bewährte Standard in der dentalen Lokalanästhesie

- [1] Symposium „20 Jahre Ultracain®“, Veranstalter: Hoechst Marion Roussel, 22. und 23. November 1996, Erbach.
- [2] Pressegespräch „Made in Höchst: 110 Jahre moderne Lokalanästhesie“, Veranstalter: Sanofi-Aventis Deutschland GmbH, 15. April 2015, Frankfurt.
- [3] Halling F. Verbrauch dentaler Lokalanästhetika in Deutschland und im internationalen Vergleich. Deutsche Zahnärztliche Zeitschrift (DZZ) (6/2015).
- [4] GfK – DDM Jahresbericht 2014.
- [5] Kämmerer PW, Palarie V, Daubländer M et al. Comparison of 4 % articaine with epinephrine (1:100,000) and without epinephrine in inferior alveolar block for tooth extraction: double-blind randomized clinical trial of anesthetic efficacy. Oral Surg Oral Med Oral Pathol Oral Radiol. 2012 Apr; 113(4):495–9.