ZMK 4/2023 (39) 194-198

Dr. Frederic Meyer, et al.

Remineralisation von Zähnen mit MIH (Molaren-Inzisiven-Hypomineralisation) mit einer Hydroxylapatit-Zahnpasta

[1] Giuca MR, Lardani L, Pasini M, Beretta M, Gallusi G, Campanella V. State-of-the-art on MIH. Part. 1 Definition and epidemiology. 2020.

[2] Weerheijm KL, Duggal M, Mejàre I et al. Judgement criteria for Molar Incisor Hypomineralisation (MIH) in epidemiologic studies: A summary of the European meeting on MIH held in Athens, 2003. Eur J Paediatr Dent. 2003; 4: 110–114.

[3] Lygidakis NA, Garot E, Somani C, Taylor GD, Rouas P, Wong FSL. Best clinical practice guidance for clinicians dealing with children presenting with molar-incisor-hypomineralisation (MIH): an updated European Academy of Paediatric Dentistry policy document. Eur Arch Paediatr Dent. 2022; 23 (1): 3-21.

[4] Denis M, Atlan A, Vennat E, Tirlet G, Attal J-P. White defects on enamel: Diagnosis and anatomopathology: Two essential factors for proper treatment (part 1). Int Orthod. 2013; 11: 139–165.

[5] Farah R, Drummond B, Swain M, Williams S. Linking the clinical presentation of molar-incisor hypomineralisation to its mineral density. Int J Paediatr Dent. 2010; 20: 353–360.

[6] Huang TTY, Jones AS, He LH, Darendeliler MA, Swain MV. Characterisation of enamel white spot lesions using X-ray micro-tomography. J Dent. 2007; 35: 737–743.

[7] Zhao D, Dong B, Yu D, Ren Q, Sun Y. The prevalence of molar incisor hypomineralisation: evidence from 70 studies. Int J Paediatr Dent. 2018; 28: 170–179.

[8] Schwendicke F, Elhennawy K, Reda S, Bekes K, Manton DJ, Krois J. Global burden of molar incisor hypomineralisation. J Dent. 2018; 68: 10–18.

[9] Garot E, Rouas P, Somani C, Taylor GD, Wong F, Lygidakis NA. An update of the aetiological factors involved in molar incisor hypomineralisation (MIH): a systematic review and meta-analysis. Eur Arch Paediatr Dent. 2022; 23 (1): 23-38.

[10] Chawla N, Messer LB, Silva M. Clinical Studies on Molar-Incisor-Hypomineralisation Part 1: Distribution and Putative Associations. Eur Arch Paediatr Dent. 2008; 9: 180–190.

[11] Rodd HD, Boissonade FM, Day PF. Pulpal Status of Hypomineralized Permanent Molars. Pediatr Dent. 2007; 29: 514–520.

[12] Jälevik B, Klingberg GA. Dental treatment, dental fear and behaviour management problems in children with severe enamel hypomineralisation of their permanent first molars. Int J Paediatr Dent. 2002; 12: 24–32.

[13] Dantas-Neta NB, Moura L de FA de D, Cruz PF et al. Impact of molar-incisor hypomineralisation on oral health-related quality of life in schoolchildren. Braz Oral Res. 2016; 30. doi:10.1590/1807-3107BOR-2016.vol30.0117.

[14] Somali C, Taylor GD, Gadot E, Rouas P, Lygidakis NA, Wong FSL. An update of treatment modalities in children and adolescents with teeth affected by molar incisor hypomineralisation (MIH): a systematic review. Eur Arch Paediatr Dent. 2022; 23 (1): 39-64.

[15] Lygidakis NA, Dimou G, Stamataki E. Retention of fissure sealants using two different methods of application in teeth with hypomineralized molars (MIH): a 4 year clinical study. Eur Arch Paediatr Dent. 2009; 10: 223–226.

[16] Ehlers V, Reuter AK, Kehl E-B et al. Efficacy of a Toothpaste Based on Microcrystalline Hydroxyapatite on Children with Hypersensitivity Caused by MIH: A Randomised Controlled Trial. Oral Health Prev Dent. 2021; 19: 647–658.

[17] Crombie FA, Cochrane NJ, Manton DJ, Palamara JEA, Reynolds EC. Mineralisation of Developmentally Hypomineralized Human Enamel in vitro. Caries Res. 2013; 47: 259–263.

[18] Olgen IC, Sonmez H, Bezgin T. Effects of different remineralization agents on MIH defects: a randomized clinical study. Clin Oral Investig. 2022; 26: 3227–3238.

[19] Olmo-González B, Moreno-López R, Ribera-Uribe M. Dental management strategies for Molar Incisor Hypomineralisation. Pediatr Dent J. 2020; 30: 139–154.

[20] Singh SK, Goyal A, Gauba K, Rathore M. A Comparative Evaluation of CPP-ACP Cream and Fluoride Varnish in Remineralization of MIH-affected Teeth Using Laser Fluorescence. Pediatr Dent. 2021; 4: 117–121.

[21] Najibfard K, Ramalingam K, Chedjieu I, Amaechi B. Remineralization of early caries by a nano-hydroxyapatite dentifrice. J Clin Dent. 2011; 22: 139–43.

[22] O'Hagan-Wong K, Enax J, Meyer F, Ganss B. The use of hydroxyapatite toothpaste to prevent dental caries. Odontology 2022; 110: 223–230.

[23] Amaechi BT, AbdulAzees PA, Alshareif DO et al. Comparative efficacy of a hydroxyapatite and a fluoride toothpaste for prevention and remineralization of dental caries in children. BDJ Open. 2019; 5: 1–9.

[24] Amaechi BT, Phillips TS, Evans V et al. The Potential of Hydroxyapatite Toothpaste to Prevent Root Caries: A pH-Cycling Study. Clin Cosmet Investig Dent. 2021; 13: 315–324.

[25] Amaechi BT, AbdulAzees PA, Okoye LO, Meyer F, Enax J. Comparison of hydroxyapatite and fluoride oral care gels for remineralization of initial caries: a pH-cycling study. BDJ Open. 2020; 6: 9.

[26] Tschoppe P, Zandim DL, Martus P, Kielbassa AM. Enamel and dentine remineralization by nano-hydroxyapatite toothpastes. J Dent. 2011; 39: 430–437.

[27] Ekambaram M, Itthagarun A, King NM. Ingestion of fluoride from dentifrices by young children and fluorosis of the teeth-a literature review. J Clin Pediatr Dent. 2011; 36: 111–122.

[28] Green R, Lanphear B, Hornung R et al. Association Between Maternal Fluoride Exposure During Pregnancy and IQ Scores in Offspring in Canada. JAMA Pediatr. 2019; 173: 940–948.

[29] Farmus L, Till C, Green R et al. Critical windows of fluoride neurotoxicity in Canadian children. Environ Res. 2021; 200: 111315.

[30] Fabritius-Vilpoux K, Enax J, Herbig M, Raabe D, Fabritius H-O. Quantitative affinity parameters of synthetic hydroxyapatite and enamel surfaces in vitro. Bioinspired Biomim Nanobiomaterials 2019; 8: 141–153.

[31] Roveri N, Battistella E, Bianchi CL et al. Surface Enamel Remineralization: Biomimetic Apatite Nanocrystals and Fluoride Ions Different Effects. J Nanomater. 2009; 2009. doi:10.1155/2009/746383.

[32] Paszynska E, Pawinska M, Gawriolek M et al. Impact of a toothpaste with microcrystalline hydroxyapatite on the occurrence of early childhood caries: a 1-year randomized clinical trial. Sci Rep. 2021; 11: 2650.

[33] Hu M-L, Zheng G, Lin H, Yang M, Zhang Y-D, Han J-M. Network meta-analysis on the effect of desensitizing toothpastes on dentine hypersensitivity. J Dent. 2019; 88: 103170.

[34] Amaechi BT, Lemke KC, Saha S, Luong MN, Gelfond J. Clinical efficacy of nanohydroxyapatite-containing toothpaste at relieving dentin hypersensitivity: an 8 weeks randomized control trial. BDJ Open. 2021; 7: 1–8.

[35] Kensche A, Holder C, Basche S, Tahan N, Hannig C, Hannig M. Efficacy of a mouthrinse based on hydroxyapatite to reduce initial bacterial colonisation in situ. Arch Oral Biol. 2017; 80: 18–26.

[36] Meyer F, Enax J. Hydroxyapatite in Oral Biofilm Management. Eur J Dent. 2019; 13: 287–290.

[37] Fabritius-Vilpoux K, Enax J, Mayweg D et al. Ultrastructural changes of bovine tooth surfaces under erosion in presence of biomimetic hydroxyapatite. Bioinspired Biomim Nanobiomaterials 2021; 10: 132–145.

[38] Lelli M, Putignano A, Marchetti M et al. Remineralization and repair of enamel surface by biomimetic Zn-carbonate hydroxyapatite containing toothpaste: a comparative in vivo study. Front Physiol. 2014; 5: 333.

[39] Epple M, Meyer F, Enax J. A critical review of modern concepts for teeth whitening. Dent J. 2019; 7: 79.

[40] Shang R, Kaisarly D, Kunzelmann K-H. Tooth whitening with an experimental toothpaste containing hydroxyapatite nanoparticles. BMC Oral Health 2022; 22: 331.

[41] Cieplik F, Rupp CM, Hirsch S et al. Ca2+ release and buffering effects of synthetic hydroxyapatite following bacterial acid challenge. BMC Oral Health 2020; 20: 1–8.

[42] Enax J, Fabritius H-O, Fabritius-Vilpoux K, Amaechi BT, Meyer F. Modes of action and clinical efficacy of particulate hydroxyapatite in preventive oral health care- state of the art. Open Dent J. 2019; 13: 274–287.

[43] Enax J, Fabritius H-O, Meyer F, Amaechi BT. Preventing oral diseases with biomimetic hydroxyapatite. J Multidiscip Care Decis Dent. 2022; 8: 9–14.

[44] Epple M. Review of potential health risks associated with nanoscopic calcium phosphate. Acta Biomater. 2018; 77: 1–14.

[45] Meyer F, Enax J, Amaechi BT et al. Hydroxyapatite as remineralization agent for children’s dental care. Front Dent Med. 2022; 3: 1-10.

[46] Limeback H, Enax J, Meyer F. Biomimetic hydroxyapatite and caries prevention: a systematic review and meta-analysis. Can J Dent Hyg. 2021; 55: 148.

[47] Kirk RE. Practical Significance: A Concept Whose Time Has Come. Educ Psychol Meas. 1996; 56: 746–759.

[48] Steyn HS. Practical significance of the difference in means. SA J Ind Psychol. 2000; 26: 1–3.

[49] Schmitz JE, Teepe JD, Hu Y, Smith CE, Fajardo RJ, Chun Y-HP. Estimating Mineral Changes in Enamel Formation by Ashing/BSE and MicroCT. J Dent Res. 2014; 93: 256–262.

[50] Sreebny LM, Valdini A. Xerostomia. A neglected symptom. Arch Intern Med. 1987; 147: 1333–1337.

[51] Elhennawy K, Manton DJ, Crombie F et al. Structural, mechanical and chemical evaluation of molar-incisor hypomineralisation-affected enamel: A systematic review. Arch Oral Biol. 2017; 83: 272–281.

[52] Tenuta LMA, Cury JA. Fluoride: its role in dentistry. Braz Oral Res. 2010; 24: 9–17.

[53] Zohoori FV, Maguire A. Are there good reasons for fluoride labelling of food and drink? Br Dent J. 2018; 224: 215–217.

[54] Zero DT. In Situ Caries Models. Adv Dent Res. 1995; 9: 214–230.

[55] Ajaj MT, Al-Khateeb SN, Al-Batayneh OB. Effect of different acid etchants on the remineralization process of white-spot lesions: An in vitro study. Am J Dent. 2020; 33: 43–47.

[56] Peariasamy K, Anderson P, Brook A h. A quantitative study of the effect of pumicing and etching on the remineralisation of enamel opacities. Int J Paediatr Dent. 2001; 11: 193–200.

[57] Crombie FA, Manton DJ, Palamara JE, Zalizniak I, Cochrane NJ, Reynolds EC. Characterisation of developmentally hypomineralized human enamel. J Dent. 2013; 41: 611–618.

[58] Farah RA, Monk BC, Swain MV, Drummond BK. Protein content of molar–incisor hypomineralisation enamel. J Dent. 2010; 38: 591–596.

[59] Elhennawy K, Schwendicke F. Managing molar-incisor hypomineralisation: A systematic review. J Dent. 2016; 55: 16–24.

[60] Gevert MV, Soares R, Wambier LM et al. How is the quality of the available evidence on molar-incisor hypomineralisation treatment? An overview of systematic reviews. Clin Oral Investig. 2022; 26 (10): 5989-6002.

[61] Schlagenhauf U, Kunzelmann K-H, Hannig C et al. Impact of a non-fluoridated microcrystalline hydroxyapatite dentifrice on enamel caries progression in highly caries-susceptible orthodontic patients: A randomized, controlled 6-month trial. J Investig Clin Dent. 2019; 10: e12399.

[62] Amaechi, B.T., Farah, R., Liu, J.A., Phillips, T.S., Perozo, B.I., Kataoka, Y., Meyer, F., Enax, J. Remineralization of molar incisor hypomineralisation (MIH) with a hydroxyapatite toothpaste: an in-situ study, BDJ Open. 2022; 8 (1), 33.

ZMK 4/2023 (39), 152-155

B. Abou-Ayash, C. Kobbe, M. Wolf, I. Knaup, H. Meyer-Lueckel, R. J. Wierichs

Bewertung des Maskierungseffekts durch Kariesinfiltration bei post-orthodontischen, kariösen Initialläsionen: 1-Jahres-Follow-up

[1] Sundararaj D, et al. Critical evaluation of incidence and prevalence of white spot lesions during fixed orthodontic appliance treatment: A meta-analysis. J Int Soc Prev Community Dent. 2015; 5 (6): 433-9.
[2] Boersma JG, et al. Caries prevalence measured with QLF after treatment with fixed orthodontic appliances: influencing factors. Caries Res. 2005; 39 (1): 41-7.
[3] Ogaard B. Prevalence of white spot lesions in 19-year-olds: a study on untreated and orthodontically treated persons 5 years after treatment. Am J Orthod Dentofacial Orthop. 1989; 96 (5): 423-7.
[4] Kamber R, et al. Efficacy of sealants and bonding materials during fixed orthodontic treatment to prevent enamel demineralization: a systematic review and meta-analysis. Sci Rep. 2021; 11 (1): 16556.
[5] Knosel M, et al. External bleaching effect on the color and luminosity of inactive white-spot lesions after fixed orthodontic appliances. Angle Orthod. 2007; 77 (4): 646-52.
[6] Bourouni S, et al. Efficacy of resin infiltration to mask post-orthodontic or non-post-orthodontic white spot lesions or fluorosis - a systematic review and meta-analysis. Clin Oral Investig. 2021; 25(8): 4711-4719.
[7] Kannan A, Padmanabhan S. Comparative evaluation of Icon(R) resin infiltration and Clinpro XT varnish on colour and fluorescence changes of white spot lesions: a randomized controlled trial. Prog Orthod. 2019; 20 (1): 23.
[8] Willmot DR. White lesions after orthodontic treatment: does low fluoride make a difference? J Orthod. 2004; 31(3): 235-42; discussion 202.
[9] Schmidlin PR, et al. Histological, morphological, profilometric and optical changes of human tooth enamel after microabrasion. Am J Dent. 2003; 16 Spec No: 4A-8A.
[10] Knosel M, Eckstein A, Helms HJ. Durability of esthetic improvement following Icon resin infiltration of multibracket-induced white spot lesions compared with no therapy over 6 months: a single-center, split-mouth, randomized clinical trial. Am J Orthod Dentofacial Orthop. 2013; 144 (1): 86-96.
[11] Kobbe C, et al. Evaluation of the value of re-wetting prior to resin infiltration of post-orthodontic caries lesions. J Dent. 2019; 91: 103243.
[12] Jansen EE, et al. Do bleaching gels affect the stability of the masking and caries-arresting effects of caries infiltration-in vitro. Clin Oral Investig. 2021; 25 (6): 4011-4021.
[13] Ceci M, et al. Resin infiltrant for non-cavitated caries lesions: evaluation of color stability. J Clin Exp Dent. 2017; 9 (2): e231-e237.
[14] Paris S, et al. Masking of white spot lesions by resin infiltration in vitro. J Dent. 2013; 41 Suppl 5: e28-34.
[15] Gu X, et al. Esthetic improvements of postorthodontic white-spot lesions treated with resin infiltration and microabrasion: A split-mouth, randomized clinical trial. Angle Orthod. 2019; 89 (3): 372-377.
[16] Knosel M, Eckstein A, Helms HJ. Long-term follow-up of camouflage effects following resin infiltration of post orthodontic white-spot lesions in vivo. Angle Orthod. 2019; 89 (1): 33-39.
[17] Knaup I, et al. Correlation of quantitative light-induced fluorescence and qualitative visual rating in infiltrated post-orthodontic white spot lesions. Eur J Orthod. 2022.

[18] Wierichs RJ, Kobbe C, Meyer-Lueckel H. Wie der Maskierungserfolg von postorthodontischen White-Spot-Läsionen vor der Infiltration abgeschätzt werden kann. Dtsch Zahnärztl Z. 2020; 75(5): 60-66.
[19] Mazur M, et al. Objective and subjective aesthetic performance of icon(R) treatment for enamel hypomineralization lesions in young adolescents: A retrospective single center study. J Dent. 2018; 68: 104-108.
[20] Andrade R, et al. Clinical evaluation of the immediate masking effect of enamel white spot lesions treated with an infiltrant resin. Int J Esthet Dent. 2020; 15 (3): 306-316.
[21] Kim S, et al. The evaluation of resin infiltration for masking labial enamel white spot lesions. Int J Paediatr Dent. 2011; 21 (4): 241-8.
[22] Holmen L, Thylstrup A, Artun J. Surface changes during the arrest of active enamel carious lesions in vivo. A scanning electron microscope study. Acta Odontol Scand. 1987; 45 (6): 383-90.
[23] Wierichs RJ, et al. Evaluation of the masking efficacy of caries infiltration in post-orthodontic initial caries lesions: 1-year follow-up. Clin Oral Investig. 2023.

Zahnmedizinische Probleme bei Kleinkindern: Karies & Co.

Dr. Julian Schmoeckel

ZMK 4/2023 (39), 142-150

1. Andersson L, Andreasen JO (2007). Soft tissue injuries. In: Andreasen JO, Andreasen FM, Andersson L: Textbook and colour atlas of traumatic injuries to the teeth. 4th ed. Blackwell Munksgaard, Kopenhagen 2007, p. 584.
2. Bauss O, Schäfer W, Sadat-Khonsari R, Knösel M (2010). Influence of orthodontic extrusion on pulpal vitality of traumatized maxillary incisors. J Endod 36:203-207.
3. Brüllmann D, Schulze R K, d'Hoedt B (2011). Therapeutisches Vorgehen bei Frontzahntraumata (Review). Deutsches Ärzteblatt 108 (34-35): 565-570.
4. Bücher K, Neumann C, Hickel R, Kühnisch J (2013). Traumatic dental injuries at a German university clinic 2004-2008. Dent Traumatol 29(2):127-33.
5. BZÄK/KZBV (Bundeszahnärztekammer/Kassenzahnärztliche Bundesvereinigung): Beck J, Eßer W, Gösling J, Husemann J, Kesler H, Oesterreich D, Santamaria Sanchez R, Schmoeckel J, Splieth C, Stein H, Ziller S. Praktischer Ratgeber für die zahnärztliche Praxis - Frühkindliche Karies vermeiden. Deutschland. Berlin 2019. https://www.bzaek.de/fileadmin/PDFs/b16/ecc-ratgeber.pdf
6. DGPZM (Deutschen Gesellschaft für Präventivzahnmedizin): Neue Empfehlungen für Kinderzahnpasten mit Fluorid. https://www.dgpzm.de/neue-empfehlungen-fuer-kinderzahnpasten-mit-fluorid
7. Deutsche Gesellschaft für Mund-, Kiefer- und Gesichtschirurgie. Therapie des dentalen Traumas bleibender Zähne, 2015. http://www.dgzmk.de/uploads/tx_szdgzmkdocuments/traumalang.pdf. Accessed on: 16.04.2018.
8. Elfrink ME, Ghanim A, Manton DJ, Weerheijm KL. Standardised studies on Molar Incisor Hypomineralisation (MIH) and Hypomineralised Second Primary Molars (HSPM): a need. Eur Arch Paediatr Dent. 2015;16(3):247-55.
9. IDZ: Fünfte Deutsche Mundgesundheitsstudie – DMS V, Köln 2016
10. IQWiG. Fluoridlackapplikation im Milchgebiss zur Verhinderung von Karies. Rapid Report N17-03. Institut für Qualität und Wirtschaftlichkeit im Gesundheitswesen (IQWiG). Version 29.03.2018. ISSN: 1864-2500.
11. Lam R (2016). Epidemiology and outcomes of traumatic dental injuries: a review of the literature. In Australian dental journal 61 Suppl 1, pp. 4–20. DOI: 10.1111/adj.12395.
12. Maurer P (2010). Epidemiologie der Verletzungen im Kiefer-Gesichtsbereich. In: Schubert J: Traumatologie für Zahnärzte. 1. Aufl. Sanofi-Aventis, Berlin 2010, pp. 10-12.
13. Miller W.,Rollnick S.: Motivierende Gesprächsführung. Lambertus, Freiburg im Breisgau 2015.
14. Public Health England (2017). Local authorities improving oral health: commissioning better oral health for children and young people. An evidence-informed toolkit for local authorities. London. https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/601833/delivering_better_oral_health_summary.pdf
15. Sälzer S, Alkilzy M, Slot DE, Dörfer CE, Schmoeckel J, Splieth CH (2017). Chairs of Working Group 3; ORCA. Socio-behavioural aspects in the prevention and control of dental caries and periodontal diseases at an individual and population level. J Clin Periodontol. Mar;44 Suppl 18:S106-S115. doi: 10.1111/jcpe.12673.

16. TEAM DAJ (2017). (R. Basner, Dr. R. M. Santamaría, Dr. J. Schmoeckel, Dr. E. Schüler, Prof. Dr. Ch. H. Splieth). Epidemiologische Begleituntersuchungen zur Gruppenprophylaxe. Deutsche Arbeitsgemeinschaft für Jugendzahnpflege e.V. (DAJ). Gutachten, Bonn.
17. Wagner Y, Heinrich-Weltzien R (2016). Evaluation of an interdisciplinary preventive programme for early childhood caries: findings of a regional German birth cohort study (2016). Clin Oral Investig 20: 1943-1952
18. Wagner Y, Knaup I, Knaup TJ, Jacobs C, Wolf M (2020). Influence of a programme for prevention of early childhood caries on early orthodontic treatment needs. Clin Oral Investig. doi: 10.1007/s00784-020-03295-4
19. Broadbent JM, Thomson WM, R Poulton R (2008). Trajectory patterns of dental caries experience in the permanent dentition to the fourth decade of life. J Dent Res Jan;87(1):69-72. doi: 10.1177/154405910808700112
20. Berg B, Cremer M, Flothkötter M, Koletzko B, Krämer N, Krawinkel M, Lawrenz B, Przyrembel H, Schiffner U, Splieth C, Vetter K, Weißenborn A. Kariesprävention im Säuglings- und frühen Kindesalter. Handlungsempfehlungen des bundesweiten Netzwerks Gesund ins Leben. Monatsschr Kinderheilkd 2021, 169. DOI: 10.1007/s00112-021-01167-z
21. Eur Arch Paediatr Dent . 2019 Dec;20(6):507-516. doi: 10.1007/s40368-019-00464-2 . Epub 2019 Nov 8.

Guidelines on the use of fluoride for caries prevention in children: an updated EAPD policy document

K J Toumba [1], S Twetman [2], C Splieth [3], C Parnell [4], C van Loveren [5], N A Lygidakis [6]

PMID: 31631242 ; DOI: 10.1007/s40368-019-00464-2