

Pekkton® ivoy – ein Hochleistungspolymer in der Implantat-, Total- und Hybridprothetik

ZT Robert Arvai, ZA Nasser Chirazi

ZMK 6 (32), S. 386-384

- Stawarczyk B, Thrun H, Eichberger M, Roos M, Edelhoff D, Schweiger J, Schmidlin PR: Effect of different surface pretreatments and adhesives on the load-bearing capacity of veneered 3-unit PEEK FDPs. *J Prosthet Dent* 114 (5), 666–673 (2015).
- Stawarczyk B, Beuer F, Wimmer T, Jahn D, Sener B, Roos M, Schmidlin PR: Polyetheretherketone-a suitable material for fixed dental prostheses? *J Biomed Mater Res B Appl Biomater* 101 (7), 1209–1216 (2013).
- Taufall S, Eichberger M, Schmidlin PR, Stawarczyk B: Fracture load and failure types of different veneered polyetheretherketone fixed dental prostheses. *Clin Oral Investig* 2016 Mar 15 [Epub ahead of print]. DOI: 10.1007/s00784-016-1777-4
- Wang X, Lu T, Wen J, Xu L, Zeng D, Wu Q, Cao L, Lin S, Liu X, Jiang X. Selective responses of human gingival fibroblasts and bacteria on carbon fiber reinforced polyetheretherketone with multilevel nanostructured TiO₂. *Biomaterials* 83, 207–218 (2016).
- Schwitalla AD, Spintig T, Kallage I, Müller WD: Flexural behavior of PEEK materials for dental application. *Dent Mater* 31 (11), 1377–1384 (2015).
- Schwitalla AD, Spintig T, Kallage I, Müller WD: Pressure behavior of different PEEK materials for dental implants. *J Mech Behav Biomed Mater* 54, 295–304 (2016).
- Sampaio M, Buciumeanu M, Henriques B, Silva FS, Souza JC, Gomes JR: Comparison between PEEK and Ti₆Al₄V concerning micro-scale abrasion wear on dental applications. *J Mech Behav Biomed Mater* 60, 212–219 (2016).
- Sturz CR, Faber FJ, Scheer M, Rothamel D, Neugebauer J: Effects of various chair-side surface treatment methods on dental restorative materials with respect to contact angles and surface roughness. *Dent Mater J* 34 (6), 796–813 (2015).
- Najeeb S, Zafar MS, Khurshid Z, Siddiqui F: Applications of polyetheretherketone (PEEK) in oral implantology and prosthodontics. *J Prosthodont Res* 60 (1), 12–19 (2016).
- Maló P, de Sousa ST, De Araújo Nobre M, Moura Guedes C, Almeida R, Roma Torres A, Legatheaux J, Silva A: Individual lithium disilicate crowns in a full-arch, implant-supported rehabilitation: a clinical report. *J Prosthodont* 23 (6), 495–500 (2014).

Publications overview of Pekkton®

1. Alsadon O., Evaluation of the Optical Properties of PEKK based Restoration. Poster 3667, IADR Boston, March 2015.
2. Alsadon O., Structural Integrity of Poly-Ether-Ketone-Ketone (PEKK) based Bi-layered Molar Crowns, Poster BSODR 2015.
3. Arvai R., Una nuova classe di materiali "in sé", Pekkton® ivory il nuovo polimero ad alte prestazioni. Un caso clinico. Quintessenza Odontotecnica 2014;10:64-72.
4. Arvai R., Das neue Hochleistungspolymer Pekkton® ivory – eine Werkstoffklasse für sich. Quintessenz Zahntech 2014;40(11):1454-1464.
5. Chirazi N.; In der Humanmedizin bestens bekannt, in der Zahnmedizin "fast" ein Neuling (Zahn-Zeitung Schweiz Nr. 3/2016)
6. Copponnex T., DeCarmin A.: Reevaluating Thermoplastics. European Medical Device Manufacturer, March/April 2009.
7. Copponnex T.: Like a chameleon. Medical Device Developments, 2010
8. Copponnex T., Blümli M.: New material approaches in dental technology. meditec, October 2011
9. Digital Dentistry Yearbook (JP) 2015
10. Fuhrer M.; In-Vitro Untersuchung der mech. Eigenschaften von Primärstegkonstruktionen aus HLP (Masterarbeit 2015)
11. Fuhrmann G., Steiner M., Freitag-Wolf S., Kern M.: Resin bonding to three types of polyaryletherketones (PAEKs) - Durability and influence of surface conditioning. Dental Materials 2014 Mar;30(3):357-63.
12. Fuhrmann G., Dissertation: Klebeverbund zu PEKK – Einfluss der Konditionierungsverfahren und Hydrolysebeständigkeit. 2014.
13. Gobert B.: C'est quoi le PEKK ? Technologie Dentaire, Décembre 2013 N° 324.
14. Gobert B.: Faux moignon anatomique en Pekkton. Technologie Dentaire, Décembre 2013 N° 324.
15. Keilig L., Katzenbach A., Weber A., Stark H., Bourauel C.: Biomechanische Untersuchung eines Hochleistungspolymer für den Ersatz in der dentalen Prothetik. Vortrag auf der Jahrestagung der Deutschen Gesellschaft für Biomechanik (DGfB) 2013 in Ulm.
16. Keilig L., Katzenbach A., Weber A., Ottersbach K., Brune M., Stark H., Bourauel C.: Dauerlastuntersuchung an Kronen aus einem Hochleistungspolymer. DGPro 2014, Aachen.
17. Keilig L., Katzenbach A., Weber A., Stark H., Bourauel C.: Dauerlastuntersuchung an Kronen aus einem Hochleistungspolymer. Poster DGPro 2014, Aachen.
18. Keilig L., Katzenbach A., Weber A., Stark H., Bourauel C.: Fatigue testing of crowns made from a high performance polymer. EAO 2014, Rom.
19. Keilig L., Katzenbach A., Weber A., Stark H., Bourauel C.: Fatigue testing of crowns made from a high performance polymer. Poster EAO 2014, Rom.
20. Keilig L., Stark H., Bourauel C.: Biomechanics of Three- and Four-Unit-Bridges Made of Different Framework Materials – A Numerical Study. EAO 2014, Rom.
21. Keilig L., Stark H., Bourauel C.: Biomechanics of Three- and Four-Unit-Bridges Made of Different Framework Materials – A Numerical Study. Poster EAO 2014, Rom.
22. Keilig L., Stark H., Bourauel C.: Numerical Analysis of Dental Bridges. Made of different Framework Materials. Presentation CMBBE 10-2014, Amsterdam.
23. Kyowa; QDT Vol.40/2015 November Page 1627 – 1635.
24. Pham V.T.: Pekkton - Nouveau polymère hautes performances. Technologie Dentaire, Mars 2014 N° 327.
25. Pham V.T.: Pekkton - A new high-performance polymer. Dental Technologies, US Edition, 2014 n°109.
26. Silla M.: Polyetherketonketon (PEKK) als Restaurationswerkstoff in der modernen Zahnmedizin: eine Literaturübersicht. Quintessenz Zahntech 2016;42(2):2–16.
27. Taner Ch.; Computer guided immediate rehabilitation of completely edentulous jaws with fixed prostheses supported by implants (Euroteknika Barcelona 2016)
28. Tannous F., Steiner M., Shahin R., Kern M.: Retentive forces and fatigue resistance of thermoplastic resin clasps. Dental Materials 2012 Mar;28(3):273-8.
29. Wichnalek, Metallfreier-herausnehmbarer-Zahnersatz. ZWR – Das Deutsche Zahnärzteblatt 2015; 124 (6):280-285

Der informierte Patient im Fokus: Partizipative Entscheidungsfindung in der Zahnmedizin

- [1] Beachamp TL, Childress JF: Principles of Biomedical Ethics. Oxford University Press, New York (1994).
- [2] Braun B (2009): Was halten Ärzte von "shared decision making" und "decision aids" und warum nutzen sie sie nicht intensiver? In: Forum Gesundheitspolitik.
- [3] Charles C, Gafni A, Whelan T: Self-reported use of shared decision-making among breast cancer specialists and perceived barriers and facilitators to implementing this approach. *Health Expect* 7, 338–348 (2004).
- [4] Charles C, Gafni A, Whelan T: Shared decision-making in the medical encounter: what does it mean? (or it takes at least two to tango). *Soc Sci Med* 44, 681–692 (1997).
- [5] Charles C, Whelan T, Gafni A: What do we mean by partnership in making decisions about treatment? *Bmj* 319, 780–782 (1999).
- [6] Coulter A, Magee H (2003): The European Patient of the Future. In: Open University Press, Maidenhead.
- [7] Dierks ML, Seidel G (2005): Gleichberechtigte Beziehungsgestaltung zwischen Ärzten und Patienten - wollen Patienten wirklich Partner sein? In: M Härter, A Loh and C Apies (eds) Gemeinsam entscheiden - erfolgreich behandeln. Neue Wege für Ärzte und Patienten im Gesundheitswesen. Deutscher Ärzteverlag, Köln, p 35–44.
- [8] Evans R, Edwards A, Brett J, Bradburn M, Watson E, Austoker J, Elwyn G: Reduction in uptake of PSA tests following decision aids: systematic review of current aids and their evaluations. *Patient Educ Couns* 58, 13–26 (2005).
- [9] Gallagher SM: Paternalism in healthcare decision making. *Ostomy Wound Manage* 44, 22, 24–25 (1998).
- [10] Geisler L (2002): Arzt-Patienten-Beziehung im Wandel - Stärkung des dialogischen Prinzips. In: Recht und Ethik der modernen Medizin. p 216–220.
- [11] Gigerenzer G, Edwards A: Simple tools for understanding risks: from innumeracy to insight. *Bmj* 327, 741–744 (2003).
- [12] Härter M: Partizipative Entscheidungsfindung (Shared Decision-Making) - ein von Patienten, Ärzten und der Gesundheitspolitik geforderter Ansatz setzt sich durch. *Z Ärztl Fortbild Qualitätssich* 98, 89–92 (2002).
- [13] Harter M, Müller H, Dirmmaier J, Donner-Banzhoff N, Bieber C, Eich W: Patient participation and shared decision making in Germany - history, agents and current transfer to practice. *Z Evid Fortbild Qual Gesundhwes* 105, 263–270 (2011).
- [14] Hirsch O, Keller H, Krones T, Donner-Banzhoff N: Acceptance of shared decision making with reference to an electronic library of decision aids (arriba-lib) and its association to decision making in patients. An evaluation study. *Implement Sci* 6, 70 (2011).
- [15] Johnson BR, Schwartz A, Goldberg J, Koerber A: A chairside aid for shared decision making in dentistry: a randomized controlled trial. *J Dent Educ* 70, 133–141 (2006).
- [16] Kriston L, Scholl I, Holzel L, Simon D, Loh A, Harter M: The 9-item Shared Decision Making Questionnaire (SDM-Q-9). Development and psychometric properties in a primary care sample. *Patient Educ Couns* 80, 94–99 (2010).
- [17] Kupke J, Wicht MJ, Stützer H, Derman SH, Lichtenstein NV, Noack MJ: Does the use of a visualised decision board by undergraduate students during shared decision-making enhance patients' knowledge and satisfaction? - A randomised controlled trial. *Eur J Dent Educ* 17, 19–25 (2013).
- [18] Kurtz S, Silverman J, Draper J: Teaching and learning communication skills in medicine. Radcliffe Publishing Ltd., Oxford (2005).

- [19] Legare F, Stacey D, Forest PG, Coutu MF: Moving SDM forward in Canada: milestones, public involvement, and barriers that remain. *Z Evid Fortbild Qual Gesundheitswes* 105, 245–253 (2011).
- [20] Loh A, Simon D, Levente K, Härter M: Patientenbeteiligung bei medizinischen Entscheidungen. Effekte der Partizipativen Entscheidungsfindung aus systematischen Reviews. *Deutsches Ärzteblatt* 104, 1483–1488 (2007).
- [21] Mccaffery KJ, Smith S, Shepherd HL, Sze M, Dhillon H, Jansen J, Juraskova I, Butow PN, Trevena L, Carey K, Tattersall MH, Barratt A: Shared decision making in Australia in 2011. *Z Evid Fortbild Qual Gesundheitswes* 105, 234–239 (2011).
- [22] Robertson K: Active listening: more than just paying attention. *Aust Fam Physician* 34, 1053–1055 (2005).
- [23] Scheibler F, Schwantes U, Kampmann M, Pfaff H: Shared decision-making. *GGW* 5, 23–31 (2005).
- [24] Streeck S: Dominanz und Kooperation in der neuropädiatrischen Sprechstunde. In: G Brünner, R Fiehler and W Kindt (eds) *Angewandte Diskursforschung Band 1: Grundlagen und Beispielanalysen*. Verlag für Gesprächsforschung, Radolfzell, p 174–196 (2002).

- [1] [Baldiessa RA](#), [Corrêa MB](#), [Schuch HS](#), [Collares K](#), [Nascimento GG](#), [Jardim PS](#), [Moraes RR](#), [Opdam NJ](#), [Demarco FF](#): Are there universal restorative composites for anterior and posterior teeth? J Dent 41, 1027–1035 (2013).
- [2] Buchalla W, Lennon AM: Die fluoreszenzunterstützte Kariesexkavation – Funktionsprinzip und Empfehlungen für den Gebrauch. Dtsch Zahnärztl Z 70, 110–120 (2015).
- [3] Da Rosa Rodolpho PA, Donassollo TA, Cenci MS, Loguércio AD, Moraes RR, Bronkhorst EM, Opdam NJ, Demarco FF: 22 year clinical evaluation of the performance of two posterior composites with different filler characteristics. Dent Mater 27, 955–963 (2011).
- [4] Ernst CP: Verschaltungstechnik - einmal anders. ZMK 28, 318–324 (2012).
- [5] Ernst CP: Schneidezähne in Form gebracht. ZMK 30, 636–647 (2014).
- [6] Ernst CP: Keramikteilkronen als Versorgungsoption von Zähnen mit Infraktionen. Dtsch Zahnärztl Z 70, 165–173 (2015).
- [7] Ernst CP: Vom Gold-Standard zur Gold-Alternative. ZMK 31 (3), 214–226 (2015).
- [8] Ernst CP: „Schicht im Schacht“: aktuelle Betrachtung zum Thema Bulk Fill versus Schichttechnik bei direkten Seitenzahn-Kompositrestaurationen. ZMK 31, 704–719 (2015).
- [9] [Fennis WM](#), [Kuijs RH](#), [Roeters FJ](#), [Creugers NH](#), [Kreulen CM](#): Randomized control trial of composite cuspal restorations: five-year results. J Dent Res 93, 36–41 (2014).
- [10] Hickel R, Ernst CP, Haller B, Hugo B, Kunzelmann KH, Merte KH, et al: Direkte Kompositrestaurationen im Seitenzahnbereich - Indikation und Lebensdauer. Gemeinsame Stellungnahme der Deutschen Gesellschaft für Zahnerhaltung (DGZ) und der Deutschen Gesellschaft für Zahn-, Mund- und Kieferheilkunde (DGZMK) aus dem Jahr 2005. Deutsch Zahnärztl Z 60, 543–545 (2005).
- [11] Hugo B: Optimale Approximalkontakte – Neue approximale Matrizen- und Aufbautechnik bei Frontzahnfüllungen. Ästhetische Zahnmedizin 3, 241–250 (2011).
- [12] Hugo B: Form- und Stellungskorrekturen bei Frontzähnen mittels Komposit. Quintessenz 53, 227–236 (2002).
- [13] Hugo B: Ästhetik mit Komposit. Quintessenz Verlags GmbH, Berlin (2008), ISBN 978-3938947-55-5.
- [14] Ilie N, Hickel R: Macro-, micro- and nano-mechanical investigations on silorane and methacrylate-based composites. Dent Mater 25, 810–8129 (2009).
- [15] Ilie N, Rencz A, Hickel R: Investigations towards nano-hybrid resin-based composites. Clin Oral Investig 17, 185–193 (2013).
- [16] Ilie & Stawarczyk B: Bulk-Fill-Komposite: neue Entwicklungen oder doch herkömmliche Komposite? ZMK 30, 90–96 (2014).
- [17] Klaiber B: Alles noninvasiv - Zahnformveränderung, Lückenschluss, Reduktion schwarzer Dreiecke. zm 96, 52–59 (2006).
- [18] Lühns AK: Diastemaschluss mittels direkter Technik im Frontzahnbereich. Deutsch Zahnärztl Z 66, 628–635 (2011).
- [19] [Lynch CD](#), [Opdam NJ](#), [Hickel R](#), [Brunton PA](#), [Gurgan S](#), [Kakaboura A](#), [Shearer AC](#), [Vanherle G](#), [Wilson NH](#): Guidance on posterior resin composites: Academy of Operative Dentistry - European Section. J Dent 42, 377–383 (2014).
- [20] [Scholtanus JD](#), [Ozcan M](#): Clinical longevity of extensive direct composite restorations in amalgam replacement: up to 3.5 years follow-up. J Dent 42, 1404–1410 (2014).

- [21] Toledano M, Ghinea R, Cardona JC, Cabello I, Yamauti M, Pérez MM, Osorio R: Digital image analysis method to assess the performance of conventional and self-limiting concepts in dentine caries removal. *J Dent* 41 Suppl 3, e31–38 (2013).

Ästhetik ist eine Korrelation zwischen Form und Funktion – Ein integriertes Konzept für kieferorthopädisch-kieferchirurgische Therapie von skelettalen Dysgnathien

- [1] Albino JE, Tedesco L: Esthetic need for orthodontic treatment. In Melsen B, editor: Current controversies in orthodontics. Chicago, Quintessence Publishing, 11–24 (1994).
- [2] Al-Riyami S, Cunningham SJ, Moles DR: Orthognathic treatment and temporomandibular disorders: a systematic review, Part 2. Signs and symptoms and meta-analyses. Am J Orthod Dentofacial Orthop 136 (5), 626.e1–16, discussion 626–627 (2009).
- [3] Bass NM: Dento-facial orthopaedics in the correction of the skeletal II malocclusion. Br J Orthod 9, 3–8 (1982).
- [4] Bell HW, Jacobs JD: Simultaneous repositioning of the maxilla, mandible, and chin. Treatment planning and analysis of the soft tissues. Am J Orthod 91, 28–50 (1986).
- [5] Bell WH, Proffit HR, White RP: Surgical correction of facial deformities. Saunderv Verlag, Baltimor (1980).
- [6] Bell WH, Schdeman GB: Correction of vertical maxillary deficiency: Stability and soft tissue changes. J Oral Surg 39, 666–670 (1981).
- [7] Berscheid E, Gangestade S: The social psychological implications of facial physical attractiveness. Clin Plast Surg 9, 289–296 (1982).
- [8] Buddgaard M, Melsen B, Terp S: Changes during and following total maxillary osteotomy (Le Fort I Procedure): a cephalometric study. Eur J Orthod 8, 21–29 (1986).
- [9] Canut J: Eine Analyse der dentofazialen Ästhetik. Inf Orthod Kieferorthop 28, 83–105 (1996).
- [10] Centenero SAH, Hernández-Alfaro F: 3D planning in orthognathic surgery: CAD/CAM surgical splints and prediction of the soft and hard tissues results – Our experience in 16 cases. J Craniomaxillofac Surg 40, 162–168 (2012).
- [11] Collins P, Epker BN: The alar basse Cinch: A technique for prevention of alar base flaring secondary to maxillary surgery. Oral Surg 53, 549–554 (1982).
- [12] Dal Pont G: L'osteotomia retromolare per la correzione della progenia, Minerva chir 18, 1138–1141 (1959).
- [13] Dal Pont G: Die retromolare Osteotomie zur Korrektur der Progenie, der Retrogenie und des Mordex apertus. Öst Z Stoma 58, 8–10 (1961).
- [14] Dryland-Vig KWL, Ellis III E: Diagnosis and treatment planning for the surgical-orthodontic Patient. Cli Plast Surg 16, 645–658 (1989).
- [15] Dann JJ, Fonseca RJ, Bell WH: Soft tissue changes associated with total maxillary advancement: a preliminary study. J Oral Surg 34, 19–23 (1976).
- [16] De Assis EA, Starck WJ, Epker B: Cephalometric analysis of profile nasal esthetics. Part III. Postoperativ changes after isolated superior repositioning. Int J Adult Orthod Orthognath 11, 279–288 (1996).
- [17] Ellis E: Condylar positioning devices for orthodontic surgery: Are they necessary. J Oral Maxillofac Syrg 52, 536–552 (1994).
- [18] Ellis E, Hinton RJ: Histologic examination of the temporomandibular joint after mandibular advancement with and without rigid fixation: An experimental investigation in adult Maccaca mulatta. J Oral Maxillofac Syrg 49, 1316 (1991).
- [19] Epker B: Esthetic maxillofacial surgery. Lea und Febiger Verlag, Philadelphia (1994).
- [20] Epker BN, Fish LC: Surgical-orthodontic correction of open-bite deformity. Am J Orthod 71, 278–299 (1977).
- [21] Epker BN, Fish LC: Surgical superior repositioning of the maxilla: What to do with the mandible? Am J Orthod 78, 164–191 (1980).

- [22] Farkas LG, Kolar JC: Anthropometry and art in the aesthetics of women's face. *Clin Plast Surg* 14, 599–615 (1987).
- [23] Flanary CM, Barnwell GM, Alexander JM: Patient perceptions of orthognathic surgery. *Am J Orthod* 88, 137–145 (1985).
- [24] Freihofer HPM: Changes in nasal profile after maxillary advancement in cleft and non-cleft patients. *J Maxillofac Surg* 5, 20–27 (1977).
- [25] Helm S, Siersbaek-Nielsen S, Skieller V, Björk A: Skelatal maturation of the hand in relation to maximum puberal growth in body height. *Danish Dental Journal* 75, 1223–1234 (1971).
- [26] Hogeman KE, Sarnäs KV, Willmar K: One-stage surgical and dental-orthopedic correction of bimaxillary and craniofacial dysostosis. *Fortschr Kiefer Gesichtsch* 18, 39–49 (1974).
- [27] Hui E, Hägg EUO, Tidman H: Soft tissue changes following maxillary osteotomies in cleft lip and palate and non-cleft Patients. *J Craniomaxillofac Surg* 22, 182–186 (1994).
- [28] Jensen AC, Sinclair PM, Wolford LM: Soft tissue changes associated with double jaw surgery. *Am J Orthod Dentofac Orthop* 101, 266–275 (1992).
- [29] Jacobson A: The influence of children's dentofacial appearance on their social attractiveness as judged by peers and lay adults. *Am J Orthod* 79, 399–415 (1981).
- [30] Joss CU, Vassalli IM: Stability after bilateral sagittal split osteotomy advancement surgery with rigid internal fixation: a systematic review. *J Oral Maxillofac Surg* 67, 301–313 (2009).
- [31] Kiyak HA, Hohl T, West RA: Psychologie changes in orthognathic surgery patients: a 24-month follow-up. *J Oral Maxillofac Surg* 42, 506–512 (1984).
- [32] Lazaridou-Terzoudi T, Kiyak HA, Moore R, Athanasiou AE, Melsen B: Long-term assessment of psychologic outcomes of orthognathic surgery. *J Oral Maxillofac Surg* 61, 545–552 (2003).
- [33] Legan HL, Burstone GJ: Soft tissue cephalometric analysis for orthognathic surgery. *J Oral Surg* 38, 744–751 (1980).
- [34] Lee DY, Bailey LJ, Proffit WR: Soft tissue changes after superior repositioning of the maxilla with Le Fort I osteotomy: 5-years follow-up. *Int J Adult Orthod Orthognath* 11, 301–311 (1996)-
- [35] McNamara JA, McDougall Jr PD, Dierks JM: Arch with development in Class II patients treated with extraoral force and functional jaw orthodontics. *Am J Orthodont* 52, 353–359 (1966).
- [36] Mensink G, Verweij JP, Frank MD, Eelco BJ, Richard van Merkesteyn JP: Bad split during bilateral sagittal split osteotomy of the mandible with separators: a retrospective study of 427 patients. *Br J Oral Maxillofac Surg* 51, 525–529 (2013).
- [37] Mensink G, Zweers A, Wolterbeek R, Dicker GG, Groot RH, van Merkesteyn RJ: Neurosensory disturbances one year after bilateral sagittal split osteotomy of the mandibula performed with separators: A multi-centre prospective study. *J Craniomaxillofac Surg* 40, 763–767 (2012).
- [38] Obwegeser H: The surgical correction of mandibular prognathism and retrognathia with consideration of genioplasty. *J Oral Surg* 10, 677–689 (1957).
- [39] Obwegeser H: The indication for surgical correction of mandibular deformity by sagittal splitting technique. *Br J Surg* 1, 157–168 (1963).
- [40] Obwegeser H, Trauner R: Zur Operationstechnik bei der Progenie und anderen Unterkieferanomalien. *Dtsch Zahn Mund Kieferheilk* 23, 1–26 (1955).
- [41] Petrovic AG, Stutzmann J: Reaktionsfähigkeit des tierischen und menschlichen Kondylenknorpels auf Zell- und Molekularebene im Lichte einer kybernetischen Auffassung des faszialen Wachstums. *Fortschr Kieferorthop* 49, 405–425 (1988).
- [42] Proothi M, Drew SJ, Sachs SA: Motivating factors for patients undergoing orthognathic surgery evaluation. *J Oral Maxillofac Surg* 68, 1555–1559 (2010).
- [43] Radney LJ, Jacobs JD: Soft tissue changes associated with surgical total maxillary intrusion. *Am J Orthodont* 80, 191–212 (1981).

- [44] Rosen HM: Lip-nasal esthetics following Le Fort I osteotomy. *Plast Reconstr Surg* 81, 171–182 (1988).
- [45] Schwarz AM: Die Röntgendiagnostik. Urban & Schwarzenberg, Wien (1958).
- [46] Scott O, Kijak HA: Treatment expectation versus outcomes among orthodontic surgery patients. *Int J Adult Orthod Orthognath Surg* 6, 247–255 (1991).
- [47] Solano-Hernandez B, Antonarakis GS, Scolozzi P, Kiliaridis S: Combined orthodontic and orthognathic surgical treatment for the correction of skeletal anterior open-bite malocclusion: a systematic review on vertical stability. *J Oral Maxillofac Surg* 71 (1), 98–109 (2013).
- [48] Watted N, Witt E: NMR study of TNJ changes following functional orthopaedic treatment using the „Würzburg approach“, European Orthodontic Society (EOS) 74th Congress (1998).
- [49] Watted N: Behandlung von Klasse-II-Dysgnathien – funktionskieferorthopädische Therapie unter besonderer Berücksichtigung der dentofazialen Ästhetik. *Kieferorthop* 13, 193–208 (1999).
- [50] Watted N, Bill J, Witt E: Therapy concept for the combined orthodontic-surgical treatment of Angle class II deformities with short face syndrome. New aspects for surgical lengthening of the lower face. *Clin Orthod Res* 3, 78–93 (2000).
- [51] Watted N, Bill J, Witt E, Reuther J: Lengthening of the lower face Angle class II patients with skeletal deep bite (short-face-syndrome) through combined orthodontic-surgical treatment. 75th Congress of the European Orthodontic Society Strasbourg, France(1999).
- [52] Watted N, Teuscher T, Wieber M: Vertikaler Gesichtsaufbau und Planung kieferorthopädisch-kieferchirurgischer Kombinationsbehandlungen unter besonderer Berücksichtigung der dentofazialen Ästhetik. *Kieferorthop* 16, 29–44 (2000).
- [53] Westermarck AH, Bystedt H, von Konow L, Sällström KO: Nasolabial morphology after Le Fort I osteotomies. *Int J Oral Maxillofac Surg* 20, 25–30 (1991).
- [54] Williamsone EH, Caves SA, Edenfield RJ, Morse PK: Cephalometric analysis: comparisons between maximum intercuspation and centric relation. *Am J Orthod* 74, 672–677 (1978).
- [55] Williamsone EH, Evans DL, Barton WA, Williams BH: The effect of bite plane use on terminal hinge axis location. *Angle Orthod* 47, 25–33 (1977).
- [56] Witt E: Möglichkeiten und Grenzen der kieferorthopädischen Behandlung Erwachsener. *Fortschr Kieferorthop* 52, 1–7 (1991).
- [57] Witt E: Behandlungskonzepte. In Miethke RR, Drescher D (Hrsg.): *Kleines Lehrbuch der Angel-Klasse II/1 unter besonderer Berücksichtigung der Behandlung*. Quintessenz, Berlin (1996).