

Unerwünschte Arzneimittelinteraktionen bei Polypharmazie vermeiden

- [1] Deutsches Ärzteblatt 109 (9), A-418/B-360/C-356 S. 20 (2012).
- [2] Vorobeychik Y, Gordin V, Mao J, Chen L: Combination therapy for neuropathic pain: a review of current evidence. CNS Drugs 25 (12), 1023–1034 (2011).
- [3] Maenza J, Flexner C: Combination antiretroviral therapy for HIV infection. Am Fam Physician 57 (11), 2789–2798 (1998).
- [4] DHHS Panel on Antiretroviral Guidelines for Adults and Adolescents: Guidelines for the use of antiretroviral agents in adults and adolescents living with HIV. <https://aidsinfo.nih.gov/contentfiles/lvguidelines/adultandadolescentgl.pdf> (letzter Zugriff: 02.03.2018).
- [5] Cascorbi I: Drug interactions: principles, examples and clinical consequences. Dtsch Arztebl Int 109 (33–34), 546–555 (2012).
- [6] Maher R Jr., Hanlon J, Hajjar E: Clinical consequences of polypharmacy in elderly. Expert Opin Drug Saf 13 (1), 57–65 (2014).
- [7] Turgeon J, Michaud V, Steffen L: The dangers of polypharmacy in elderly patients. JAMA Intern Med 177 (10), 1544 (2017).
- [8] Moßhammer D, Haumann H, Mörike K, Joos S: Polypharmazie – Tendenz steigend, Folgen schwer kalkulierbar. Dtsch Arztebl Int 113 (38), 627–633 (2016).
- [9] Scheidt-Nave C, Richter S, Fuchs J, Kuhlmeier A: Robert Koch-Institut, Berlin. Herausforderungen an die Gesundheitsforschung für eine alternde Gesellschaft am Beispiel „Multimorbidität“. Heidelberg: Springer Verlag (2010).
- [10] DZW. Moderne Lokalanästhetika in der Praxis: Einsatz und Wirkung. <https://www.dzw.de/index.php/moderne-lokalanaesthetika-der-praxis-einsatz-und-wirkung> (letzter Zugriff: 02.03.2018).
- [11] Onmeda 2013: Wirkstoff: Epinephrin. <https://www.onmeda.de/Wirkstoffe/Epinephrin> (letzter Zugriff: 02.03.2018).
- [12] Halling F: Aktuelle Analyse zahnärztlicher Analgetikaverordnungen. ZWR Das Deutsche Zahnärzteblatt 120 (5), 216–227 (2011).
- [13] Onmeda 2010: Wirkstoff: Acetylsalicylsäure. <https://www.onmeda.de/Wirkstoffe/Acetylsalicylsäure.html> (letzter Zugriff: 02.03.2018).
- [14] Onmeda 2013: Wirkstoff: Coffein. <https://www.onmeda.de/Wirkstoffe/Coffein.html> (letzter Zugriff: 02.03.2018).
- [15] Onmeda 2015: Wirkstoff: Codein. <https://www.onmeda.de/Wirkstoffe/Codein.html> (letzter Zugriff: 02.03.2018).

Kommentar [AK1]: Autor? Onmeda ist ein Webseitenname

Kommentar [AK2]: s.o.

Kommentar [AK3]: s.o.

Kommentar [AK4]: s.o.

[16] Onmeda 2014: Wirkstoff: Paracetamol.
<https://www.onmeda.de/Wirkstoffe/Paracetamol.html> (letzter Zugriff: 02.03.2018).

Kommentar [AK5]: S. O.

[17] Moore N, Pollack C, Butkerait P: Adverse drug reactions and drug-drug interactions with over-the-counter NSAIDs. Ther Clin Risk Manag 11, 1061–1075 (2015).

[18] Onmeda 2017: Wirkstoff: Ibuprofen.
[https://www.onmeda.de/Wirkstoffe/Ibuprofen+\(zur+Einnahme\).html](https://www.onmeda.de/Wirkstoffe/Ibuprofen+(zur+Einnahme).html) (letzter Zugriff: 02.03.2018).

Kommentar [AK6]: S. O.

[19] Sanofi-Aventis Deutschland GmbH: Fachinformation "Novalgin® Filmtabletten". Stand: März 2017.

[20] Bundesgesundheitsministerium, e-Health Gesetz.
<https://www.bundesgesundheitsministerium.de/ministerium/meldungen/2016/medikationsplan.html> Stand: 30. September 2016.

[21] Autor(en): Top 20 der meistverkauften Arzneimittel in Deutschland im Jahr 2015 (in Millionen Packungen). <https://de.statista.com/statistik/daten/studie/548361/umfrage/top-20-der-meistverkauften-arzneimittel-in-deutschland/> (letzter Zugriff: Datum).

[22] Sanofi-Aventis. Anamnesebogen.
<https://dental.sanofi.de/bestellcenter/details?item=023fc6e7-0b3a-4a17-936b-fc64833298dd> (letzter Zugriff: 02.03.2018).

[23] § 630e – Bürgerliches Gesetzbuch (BGB) neugefasst durch B. v. 02.01.2002 BGBl. I S. 42, 2909; 2003, 738; zuletzt geändert durch Artikel 1 G. v. 20.07.2017 BGBl. I S. 2787
Geltung ab 01.01.1964; FNA: 400-2 Bürgerliches Gesetzbuch, Einführungsgesetz und zugehörige Gesetze.

Pflichttext Ultracain_D-S_D-S_forte_D_ohne_Adrenalin_Fachkreise
Pflichttext Novalgin_Fachkreise

Kommentar [AK7]: ??

- [1] AAPD: Policy on the use of dental bleaching for child and adolescent patients. *Pediatr Dent* 39 (6), 90–92 (2017).
- [2] Giniger M, Macdonald J, Ziemba S, Felix H: The clinical performance of professionally dispensed bleaching gel with added amorphous calcium phosphate. *J Am Dent Assoc* 136 (3), 383–392 (2005).
- [3] He LB, Shao MY, Tan K, Xu X, Li JY: The effects of light on bleaching and tooth sensitivity during in-office vital bleaching: a systematic review and meta-analysis. *J Dent* 40 (8), 644–653 (2012).
- [4] Karadas M, Seven N: The effect of different drinks on tooth color after home bleaching. *Eur J Dent* 8 (2), 249–253 (2014).
- [5] Kina JF, Huck C, Riehl H, Martinez TC, Sacono NT, Ribeiro AP, Costa CA: Response of human pulps after professionally applied vital tooth bleaching. *Int Endod J* 43 (7), 572–580 (2010).
- [6] Noack MJ, Padrok Ö, Höfer K: Zahnaufhellung – medizinisch indiziert. *ZWR* 124 (3), 96–100 (2015).
- [7] Parinitha MS, Annapoorna BM, Tejaswi S, Shetty S, Sowmya HK: Effect of power bleaching on the fluorosis stained anterior teeth case series. *J Clin Diagn Res* 8 (8), ZJ01–03 (2014).
- [8] Polydorou O, Scheitza S, Spraul M, Vach K, Hellwig E: The effect of long-term use of tooth bleaching products on the human enamel surface. *Odontology* 106 (1), 64–72 (2018).
- [9] Tavares M, Stultz J, Newman M, Smith V, Kent R, Carpino E, Goodson JM: Light augments tooth whitening with peroxide. *J Am Dent Assoc* 134 (2), 167–175 (2003).

ZMK 34(5) 2018, S. 400-405

Dr. Markus Th. Firla

**Das ZahnAbnutzungsBewertungsSystem (ABS) zur Dokumentation
nichtkariöser Zahnhartsubstanzschäden**

- [1] Alt KW et al.: Okklusion – Kultur versus Natur. Zahnärztl Mitt 107 (10), 58–64 (2017).
- [2] Attin T, Wegehaupt FJ: Methods for assessment of dental erosion. Monogr Oral Sci 25, 123–142 (2014).
- [3] Azzopardi A et al.: A literature review of the technique to measure tooth wear and erosion. Eur J Prosthodont Restor Dent 8, 93–97 (2000).
- [4] Barfuß A: Update Erosionen. Dent Mag 29 (6), 52–59 (2001).
- [5] Bartlett DW et al.: Basic Erosive Wear Examination (BEWE): A new scoring system for scientific and clinical needs. Clin Oral Investig 12 (1), 65–68 (2008).
- [6] Bartlett DW et al.: Prevalance of tooth wear on buccal and lingual surfaces and possible risk factors in young European adults. J Dent 41 (11), 1007–1013 (2013).
- [7] Dawson PE: Functional occlusion from TMJ to smile design. Mosby Inc., St. Louis, USA (2007).
- [8] Firla MTh: Der erosive Zahnhartsubstanzverlust – eine unterschätzte Gefahr. ZMK 26 (19), 629–633 (2010).
- [9] Firla MTh: Dentale Erosionen. Zahn Prax 14 (1), 56–59 (2011).
- [10] Firla MTh: Dentale Erosionen – Nicht selten übersehen und als Gefahr noch oft unterschätzt. Dental Barometer 6 (7), 34–38 (2011).
- [11] Firla MTh: Tatort Mundhöhle – Spurensuche und Prävention erosiver Zahnschäden. Dental Barometer 8 (6), 60–63 (2013).
- [12] Guggenbichler N: Menschen im Stress – Zur Psychosomatik des Zähneknirschens. VAS Verlag, Bad Homburg (2012).
- [13] Hancocks S (Hrsg.): Zahnhartsubstanzverlust – Neue Erkenntnisse zur Erosion und Abrasion. Sonderdruck Int Dent J (4), 259–294 (2005).
- [14] Hetz G: Erosionen und andere spezielle Defekte. Dental Barometer 5 (2), 22–26 (2010).
- [15] Kielbassa AM: Erosion und Abrasion – ein lebenslanges Problem. Dent Tribune German Edition 9 (6), 4 (2012).
- [16] Kühnisch J et al.: Strukturstörungen des Zahnschmelzes und des Dentins. Teil 2: Klinisches Erscheinungsbild. Quintessenz 62 (2), 171–186 (2011).
- [17] Lobbezoo F, Naeije M: A reliability study of clinical tooth wear measurements. J Prosthet Dent 86, 597–602 (2001).

[18] Lussi A, Jaeggi T: Dentale Erosionen – Von der Diagnose zur Therapie. Quintessenz Verlag, Berlin (2009).

[19] Magne P et al.: Anatomic crown width/length ratios of unworn and worn maxillary teeth in white subjects. J Prosthet Dent 89, 453–461 (2003).

[20] Sterling C: Jung und sportlich aktiv – Diagnose: CMD. Dental Barometer 9 (2), 16–18 (2016).

[21] The Journal of Clinical Dentistry/The International Journal of Applied Dental Research: Toumba J: Tooth surface loss – a challenge to oral health. Barbour M, Rees G: The role of erosion, abrasion and attrition in tooth wear. Newby C et al.: Surface microhardness changes, enamel fluoride uptake, and fluoride availability from commercial toothpastes. Fowler C: In vitro microhardness studies on new anti-erosion desensitizing toothpaste. Creeth JE et al.: The stain removal performance of a new anti-hypersensitivity dentrifice. Zero DT et al.: Evaluation of a desensitizing test dentrifice using an in situ erosion remineralization model. 27 (4), Special Issue 85–116 (2006).

Kommentar [AK1]: Ist das so richtig?
Das sind doch mehrere Literaturstellen.

[22] Vailati F, Belser UCh: Classification and treatment of the anterior maxillary dentition affected by dental erosion: The ACE Classification. Int J Periodontics 30 (6), 559–572 (2010).

[23] Wetselaar P et al.: Reliability of an occlusal and non-occlusal tooth wear grading system – Clinical use versus dental cast assessment. Int J Prosthodont 22, 388–390 (2009).

[24] Wetselaar P, Lobbezoo F: Diagnostik der Zahnabnutzung. AkkreDidakt/Prelum, Berlin (2017).

[25] Wöhrle W: Dentale Erosionen. In: Fuhrmann A (Hrsg.): Die zahnmedizinische Fachangestellte. 7/7.2 – 7/7.2.8. Spitta Verlag, Balingen (2009).

- [1] Augusto MG, Torres C, Pucci CR, Schlueter N, Borges AB: Bond stability of an universal adhesive system to eroded/abraded dentin after deproteinization. *Oper Dent* 43, 291–300 (2018).
- [2] Azzopardi A, Bartlett DW, Watson TF, Sherriff M: The measurement and prevention of erosion and abrasion. *J Dent* 29, 395–400 (2001).
- [3] Bartlett D, Sundaram G, Moazzez R: Trial of protective effect of fissure sealants, in vivo, on the palatal surfaces of anterior teeth, in patients suffering from erosion. *J Dent* 39, 26–29 (2011).
- [4] Carvalho TS, Colon P, Ganss C, Huysmans MC, Lussi A, Schlueter N et al.: Consensus Report of the European Federation of Conservative Dentistry: Erosive tooth wear diagnosis and management. *Swiss Dent J* 126, 342–346 (2016).
- [5] Deari S, Wegehaupt FJ, Taubock TT, Attin T: Influence of different pretreatments on the microtensile bond strength to eroded dentin. *J Adhes Dent* 19, 147–155 (2017).
- [6] Flury S, Koch T, Peutzfeldt A, Lussi A, Ganss C: The effect of a tin-containing fluoride mouth rinse on the bond between resin composite and erosively demineralised dentin. *Clin Oral Investig* 17, 217–225 (2013).
- [7] Ganss C, Hardt M, Lussi A, Cocks AK, Klimek J, Schlueter N: Mechanism of action of tin-containing fluoride solutions as anti-erosive agents in dentine – an in vitro tin-uptake, tissue loss, and scanning electron microscopy study. *Eur J Oral Sci* 118, 376–384 (2010).
- [8] Ganss C, Lussi A, Peutzfeldt A, Naguib Attia N, Schlueter N: Impact of Sn/F pre-treatments on the durability of protective coatings against dentine erosion/abrasion. *PLoS One* 10, e0123889 (2015).
- [9] Ganss C, Neutard L, von Hinckeldey J, Klimek J, Schlueter N: Efficacy of a tin/fluoride rinse: a randomized in situ trial on erosion. *J Dent Res* 89, 1214–1218 (2010).
- [10] Hirce JD, Sather AH, Chao EY: The effect of topical fluorides, after acid etching of enamel, on the bond strength of directly bonded orthodontic brackets. *Am J Orthod* 78, 444–452 (1980).
- [11] Huysmans MC, Young A, Ganss C: The role of fluoride in erosion therapy. *Monogr Oral Sci* 25, 230–243 (2014).
- [12] Imamura N, Kawaguchi T, Shimizu H, Takahashi Y: Effect of three metal priming agents on the bond strength of adhesive resin cement to Ag-Zn-Sn-In alloy and component metals. *Dent Mater J* 37, 301–307 (2018).
- [13] Loomans B, Opdam N, Attin T, Bartlett D, Edelhoff D, Frankenberger R et al.: Severe tooth wear: European Consensus Statement on Management Guidelines. *J Adhes Dent* 19, 111–119 (2017).
- [14] Low T, Von Fraunhofer JA, Winter GB: The bonding of a polymeric fissure sealant to topical fluoride-treated teeth. *J Oral Rehabil* 2, 303–307 (1975).
- [15] Low T, von Fraunhofer JA, Winter GB: Influence of the topical application of fluoride on the in vitro adhesion of fissure sealants. *J Dent Res* 56, 17–20 (1977).
- [16] Oliveira B, Ulbaldini A, Sato F, Baesso ML, Bento AC, Andrade L et al.: Chemical interaction analysis of an adhesivecontaining 10-methacryloyloxydecyl dihydrogen phosphate (10-MDP) with the dentin in noncarious cervical lesions. *Oper Dent* 42, 357–366 (2017).
- [17] Petzold M: The influence of different fluoride compounds and treatment conditions on

dental enamel: a descriptive in vitro study of the CaF₂ precipitation and microstructure. *Caries Res* 35 (Suppl 1), 45–51 (2001).

[18] Peumans M, De Munck J, Van Landuyt K, Van Meerbeek B: Thirteen-year randomized controlled clinical trial of a two-step self-etch adhesive in non-carious cervical lesions. *Dent Mater* 31, 308–314 (2015).

[19] Peutzfeldt A, Jaeggi T, Lussi A: Restorative therapy of erosive lesions. *Monogr Oral Sci* 25, 253–261 (2014).

[20] Peutzfeldt A, Koch T, Ganss C, Flury S, Lussi A: Effect of tin-chloride pretreatment on bond strength of two adhesive systems to dentin. *Clin Oral Investig* 18, 535–543 (2014).

[21] Saxegaard E, Rolla G: Fluoride acquisition on and in human enamel during topical application in vitro. *Scand J Dent Res* 96, 523–535 (1988).

[22] Schlueter N, Hardt M, Lussi A, Engelmann F, Klimek J, Ganss C: Tin-containing fluoride solutions as anti-erosive agents in enamel: an in vitro tin-uptake, tissue-loss, and scanning electron micrograph study. *Eur J Oral Sci* 117, 427–434 (2009).

[23] Schlueter N, Peutzfeldt A, Ganss C, Lussi A: Does tin pre-treatment enhance the bond strength of adhesive systems to enamel? *J Dent* 41, 642–652 (2013).

[24] Schlüter N, Gressbach C, Ziemann C, Winterfeld T, Ganß C: Kompositbasierte Beschichtungen zur Prävention von Dentinerosionen. *Dtsch Zahnärztl Z* 69, 564–572 (2014).

[25] Siqueira F, Cardenas A, Gomes GM, Chibinski AC, Gomes O, Bandeca MC et al.: Three-year effects of deproteinization on the in vitro durability of resin/dentin-eroded interfaces. *Oper Dent* 43, 60–70 (2018).

[26] Umwali A, Askar H, Paris S, Schwendicke F: Radiographic, antibacterial and bond-strength effects of radiopaque caries tagging. *Sci Rep* 6, 27319 (2016).

[27] Yetkiner E, Wegehaupt FJ, Attin R, Wiegand A, Attin T: Stability of two resin combinations used as sealants against toothbrush abrasion and acid challenge in vitro. *Acta Odontol Scand* 72, 825–830 (2014).

[28] Zimmerli B, De Munck J, Lussi A, Lambrechts P, Van Meerbeek B: Long-term bonding to eroded dentin requires superficial bur preparation. *Clin Oral Investig* 16, 1451–1461 (2012).

- [1] Peppard PE, Young T, Barnet JH, Palta M, Hagen EW, Hla KM: Increased prevalence of sleep-disordered breathing in adults. *Am J Epidemiol* 177, 1006–1014 (2013).
- [2] Young T, Palta M, Dempsey J, Skatrud J, Weber S, Badr S: The occurrence of sleep-disordered breathing among middle-aged adults. *N Engl J Med* 328, 1230–1235 (1993).
- [3] Punjabi NM: The epidemiology of adult obstructive sleep apnea. *Proc Am Thorac Soc* 5, 136–143 (2008).
- [4] Young T, Peppard PE, Gottlieb DJ: Epidemiology of obstructive sleep apnea: a population health perspective. *Am J Respir Crit Care Med* 165, 1217–1239 (2002).
- [5] Sforza E, Roche F, Thomas-Anterion C, Kerleroux J, Beauchet O, Celle S et al.: Cognitive function and sleep related breathing disorders in a healthy elderly population: the SYNAPSE study. *Sleep* 33, 515–521 (2010).
- [6] Capistrano A, Cordeiro A, Capelozza Filho L, Almeida VC, Silva PI, Martinez S et al.: Facial morphology and obstructive sleep apnea. *Dental Press J Orthod* 20, 60–67 (2015).
- [7] Denolf PL, Vanderveken OM, Marklund ME, Braem MJ: The status of cephalometry in the prediction of non-CPAP treatment outcome in obstructive sleep apnea patients. *Sleep Med Rev* 27, 56–73 (2016).
- [8] McNicholas WT: Diagnosis of obstructive sleep apnea in adults. *Proc Am Thorac Soc* 5, 154–160 (2008).
- [9] Ward Flemons W, McNicholas WT: Clinical prediction of the sleep apnea syndrome. *Sleep Med Rev* 1, 19–32 (1997).
- [10] Somers VK, White DP, Amin R, Abraham WT, Costa F, Culebras A et al.: Sleep apnea and cardiovascular disease: an American Heart Association/American College of Cardiology Foundation Scientific Statement from the American Heart Association Council for High Blood Pressure Research Professional Education Committee, Council on Clinical Cardiology, Stroke Council, and Council on Cardiovascular Nursing. *J Am Coll Cardiol* 52, 686–717 (2008).
- [11] Dong JY, Zhang YH, Qin LQ: Obstructive sleep apnea and cardiovascular risk: meta-analysis of prospective cohort studies. *Atherosclerosis* 229, 489–495 (2013).
- [12] Gami AS, Hodge DO, Herges RM, Olson EJ, Nykodym J, Kara T et al.: Obstructive sleep apnea, obesity, and the risk of incident atrial fibrillation. *J Am Coll Cardiol* 49, 565–571 (2007).
- [13] Gami AS, Pressman G, Caples SM, Kanagala R, Gard JJ, Davison DE et al.: Association of atrial fibrillation and obstructive sleep apnea. *Circulation* 110, 364–367 (2004).
- [14] Verse T, Dreher A, Heiser C, Herzog M, Maurer J, Pirsig W et al.: HNO-spezifische Therapie der obstruktiven Schlafapnoe bei Erwachsenen. AWMF-Register Nr. 017/069 (2015).
- [15] Geert M et al.: S3-Leitlinie Nicht erholsamer Schlaf/Schlafstörungen – Kapitel „Schlafbezogene Atmungsstörungen“. AWMF Registernummer 063/001 (2016).
- [16] Alves M, Jr., Franzotti ES, Baratieri C, Nunes LK, Nojima LI, Ruellas AC: Evaluation of pharyngeal airway space amongst different skeletal patterns. *Int J Oral Maxillofac Surg* 41, 814–819 (2012).
- [17] Abe-Nickler MD, Portner S, Sieg P, Hakim SG: No correlation between two-dimensional measurements and three-dimensional configuration of the pharyngeal upper airway space in cone-beam computed tomography. *J Craniomaxillofac Surg* 45, 371–376 (2017).
- [18] Abrahamyan L, Sahakyan Y, Chung S, Pechlivanoglou P, Bielecki J, Carcone SM et al.: Diagnostic accuracy of level IV portable sleep monitors versus polysomnography for obstructive sleep apnea: a systematic review and meta-analysis. *Sleep Breath* (2018).

- [19] Danker-Hopfe H, Kunz D, Gruber G, Klosch G, Lorenzo JL, Himanen SL et al.: Interrater reliability between scorers from eight European sleep laboratories in subjects with different sleep disorders. *J Sleep Res* 13, 63–69 (2004).
- [20] Anandam A, Akinnusi M, Kufel T, Porhomayon J, El-Solh AA: Effects of dietary weight loss on obstructive sleep apnea: a meta-analysis. *Sleep Breath* 17, 227–234 (2013).
- [21] Benedict MA, Arterburn D: Worksite-based weight loss programs: a systematic review of recent literature. *Am J Health Promot* 22, 408–416 (2008).
- [22] Buchwald H, Avidor Y, Braunwald E, Jensen MD, Pories W, Fahrenbach K et al.: Bariatric surgery: a systematic review and meta-analysis. *Jama* 292, 1724–1737 (2004).
- [23] Greenburg DL, Lettieri CJ, Eliasson AH: Effects of surgical weight loss on measures of obstructive sleep apnea: a meta-analysis. *Am J Med* 122, 535–542 (2009).
- [24] Sarkhosh K, Switzer NJ, El-Hadi M, Birch DW, Shi X, Karmali S: The impact of bariatric surgery on obstructive sleep apnea: a systematic review. *Obes Surg* 23, 414–423 (2013).
- [25] Permut I, Diaz-Abad M, Chatila W, Crocetti J, Gaughan JP, D'Alonzo GE et al.: Comparison of positional therapy to CPAP in patients with positional obstructive sleep apnea. *J Clin Sleep Med* 6, 238–243 (2010).
- [26] Skinner MA, Kingshott RN, Filsell S, Taylor DR: Efficacy of the 'tennis ball technique' versus nCPAP in the management of position-dependent obstructive sleep apnoea syndrome. *Respirology* 13, 708–715 (2008).
- [27] Ayas NT, FitzGerald JM, Fleetham JA, White DP, Schulzer M, Ryan CF et al.: Cost-effectiveness of continuous positive airway pressure therapy for moderate to severe obstructive sleep apnea/hypopnea. *Arch Intern Med* 166, 977–984 (2006).
- [28] Kushida CA, Littner MR, Hirshkowitz M, Morgenthaler TI, Alessi CA, Bailey D et al.: Practice parameters for the use of continuous and bilevel positive airway pressure devices to treat adult patients with sleep-related breathing disorders. *Sleep* 29, 375–380 (2006).
- [29] Randerath WJ, Hein H, Arzt M, Galetke W, Nilius G, Penzel T et al.: [Consensus paper on the diagnosis and treatment of sleep disordered breathing]. *Pneumologie* 68, 106–123 (2014).
- [30] Chai-Coetzer CL, Antic NA, Rowland LS, Reed RL, Esterman A, Catcheside PG et al.: Primary care vs specialist sleep center management of obstructive sleep apnea and daytime sleepiness and quality of life: a randomized trial. *Jama* 309, 997–1004 (2013).
- [31] Rosen CL, Auckley D, Benca R, Foldvary-Schaefer N, Iber C, Kapur V et al.: A multisite randomized trial of portable sleep studies and positive airway pressure autotitration versus laboratory-based polysomnography for the diagnosis and treatment of obstructive sleep apnea: the HomePAP study. *Sleep* 35, 757–767 (2012).
- [32] Gay P, Weaver T, Loubé D, Iber C: Evaluation of positive airway pressure treatment for sleep related breathing disorders in adults. *Sleep* 29, 381–401 (2006).
- [33] Giles TL, Lasserson TJ, Smith BH, White J, Wright J, Cates CJ: Continuous positive airways pressure for obstructive sleep apnoea in adults. *Cochrane Database Syst Rev*, Cd001106 (2006).
- [34] Sanders MH, Montserrat JM, Farre R, Givelber RJ: Positive pressure therapy: a perspective on evidence-based outcomes and methods of application. *Proc Am Thorac Soc* 5, 161–172 (2008).
- [35] Madbouly EM, Nadeem R, Nida M, Molnar J, Aggarwal S, Loomba R: The role of severity of obstructive sleep apnea measured by apnea-hypopnea index in predicting compliance with pressure therapy, a meta-analysis. *Am J Ther* 21, 260–264 (2014).
- [36] Wozniak DR, Lasserson TJ, Smith I: Educational, supportive and behavioural interventions to improve usage of continuous positive airway pressure machines in adults with obstructive sleep apnoea. *Cochrane Database Syst Rev*, Cd007736 (2014).
- [37] Kushida CA, Chediak A, Berry RB, Brown LK, Gozal D, Iber C et al.: Clinical guidelines for the manual titration of positive airway pressure in patients with obstructive sleep apnea. *J Clin Sleep Med* 4, 157–171 (2008).

- [38] Levy P, Kohler M, McNicholas WT, Barbe F, McEvoy RD, Somers VK et al.: Obstructive sleep apnoea syndrome. *Nat Rev Dis Primers* 1, 15015 (2015).
- [39] Weaver TE, Grunstein RR: Adherence to continuous positive airway pressure therapy: the challenge to effective treatment. *Proc Am Thorac Soc* 5, 173–178 (2008).
- [40] Epstein LJ, Kristo D, Strollo PJ, Jr., Friedman N, Malhotra A, Patil SP et al.: Clinical guideline for the evaluation, management and long-term care of obstructive sleep apnea in adults. *J Clin Sleep Med* 5, 263–276 (2009).
- [41] Li W, Xiao L, Hu J: The comparison of CPAP and oral appliances in treatment of patients with OSA: a systematic review and meta-analysis. *Respir Care* 58, 1184–1195 (2013).
- [42] Phillips CL, Grunstein RR, Darendeliler MA, Mihailidou AS, Srinivasan VK, Yee BJ et al.: Health outcomes of continuous positive airway pressure versus oral appliance treatment for obstructive sleep apnea: a randomized controlled trial. *Am J Respir Crit Care Med* 187, 879–887 (2013).
- [43] Randerath WJ, Verbraecken J, Andreas S, Bettge G, Boudewyns A, Hamans E et al.: Non-CPAP therapies in obstructive sleep apnoea. *Eur Respir J* 37, 1000–1028 (2011).
- [44] Ngiam J, Balasubramaniam R, Darendeliler MA, Cheng AT, Waters K, Sullivan CE: Clinical guidelines for oral appliance therapy in the treatment of snoring and obstructive sleep apnoea. *Aust Dent J* 58, 408–419 (2013).
- [45] Qaseem A, Holty JE, Owens DK, Dallas P, Starkey M, Shekelle P: Management of obstructive sleep apnea in adults: A clinical practice guideline from the American College of Physicians. *Ann Intern Med* 159, 471–483 (2013).
- [46] Puhan MA, Suarez A, Lo Cascio C, Zahn A, Heitz M, Braendli O: Didgeridoo playing as alternative treatment for obstructive sleep apnoea syndrome: randomised controlled trial. *Bmj* 332, 266–270 (2006).
- [47] Camacho M, Certal V, Abdullatif J, Zaghi S, Ruoff CM, Capasso R et al.: Myofunctional therapy to treat obstructive sleep apnea: a systematic review and meta-analysis. *Sleep* 38, 669–675 (2015).
- [48] Guimaraes KC, Drager LF, Genta PR, Marcondes BF, Lorenzi-Filho G: Effects of oropharyngeal exercises on patients with moderate obstructive sleep apnea syndrome. *Am J Respir Crit Care Med* 179, 962–966 (2009).
- [49] Randerath WJ, Galetke W, Domanski U, Weitkunat R, Rühle KH: Tongue-muscle training by intraoral electrical neurostimulation in patients with obstructive sleep apnea. *Sleep* 27, 254–259 (2004).
- [50] Camacho M, Certal V, Brietzke SE, Holty JE, Guilleminault C, Capasso R: Tracheostomy as treatment for adult obstructive sleep apnea: a systematic review and meta-analysis. *Laryngoscope* 124, 803–811 (2014).
- [51] Browaldh N, Nerfeldt P, Lysdahl M, Bring J, Friberg D: SKUP3 randomised controlled trial: polysomnographic results after uvulopalatopharyngoplasty in selected patients with obstructive sleep apnoea. *Thorax* 68, 846–853 (2013).
- [52] Li HY, Wang PC, Lee LA, Chen NH, Fang TJ: Prediction of uvulopalatopharyngoplasty outcome: anatomy-based staging system versus severity-based staging system. *Sleep* 29, 1537–1541 (2006).
- [53] Marti S, Sampol G, Munoz X, Torres F, Roca A, Lloberes P et al.: Mortality in severe sleep apnoea/hypopnoea syndrome patients: impact of treatment. *Eur Respir J* 20, 1511–1518 (2002).
- [54] Goktas O, Solmaz M, Goktas G, Olze H: Long-term results in obstructive sleep apnea syndrome (OSAS) after laser-assisted uvulopalatoplasty (LAUP). *PLoS One* 9, e100211 (2014).
- [55] Sundaram S, Bridgman SA, Lim J, Lasserson TJ: Surgery for obstructive sleep apnoea. *Cochrane Database Syst Rev*, Cd001004 (2005).
- [56] Won CH, Li KK, Guilleminault C: Surgical treatment of obstructive sleep apnea: upper airway and maxillomandibular surgery. *Proc Am Thorac Soc* 5, 193–199 (2008).

- [57] Choi JH, Kim SN, Cho JH: Efficacy of the Pillar implant in the treatment of snoring and mild-to-moderate obstructive sleep apnea: a meta-analysis. *Laryngoscope* 123, 269–276 (2013).
- [58] Maurer JT, Sommer JU, Hein G, Hormann K, Heiser C, Stuck BA: Palatal implants in the treatment of obstructive sleep apnea: a randomised, placebo-controlled single-centre trial. *Eur Arch Otorhinolaryngol* 269, 1851–1856 (2012).
- [59] Back LJ, Hytonen ML, Roine RP, Malmivaara AO: Radiofrequency ablation treatment of soft palate for patients with snoring: a systematic review of effectiveness and adverse effects. *Laryngoscope* 119, 1241–1250 (2009).
- [60] Back LJ, Liukko T, Rantanen I, Peltola JS, Partinen M, Ylikoski J et al.: Radiofrequency surgery of the soft palate in the treatment of mild obstructive sleep apnea is not effective as a single-stage procedure: A randomized single-blinded placebo-controlled trial. *Laryngoscope* 119, 1621–1627 (2009).
- [61] Farrar J, Ryan J, Oliver E, Gillespie MB: Radiofrequency ablation for the treatment of obstructive sleep apnea: a meta-analysis. *Laryngoscope* 118, 1878–1883 (2008).
- [62] Strollo PJ, Jr., Soose RJ, Maurer JT, de Vries N, Cornelius J, Froymovich O et al.: Upper-airway stimulation for obstructive sleep apnea. *N Engl J Med* 370, 139–149 (2014).
- [63] Woodson BT, Gillespie MB, Soose RJ, Maurer JT, de Vries N, Steward DL et al.: Randomized controlled withdrawal study of upper airway stimulation on OSA: short- and long-term effect. *Otolaryngol Head Neck Surg* 151, 880–887 (2014).
- [64] Bangiyev JN, Traboulsi H, Abdulhamid I, Rozzelle A, Thottam PJ: Sleep architecture in Pierre-Robin sequence: The effect of mandibular distraction osteogenesis. *Int J Pediatr Otorhinolaryngol* 89, 72–75 (2016).
- [65] Vezina JP, Blumen M, Buchet I, Chabolle F: Sleep-disordered breathing: choosing the right cephalometric analysis. *J Oral Maxillofac Surg* 70, 1442–1448 (2012).
- [66] Pirklbauer K, Russmueller G, Stiebellehner L, Nell C, Sinko K, Millesi G et al.: Maxillomandibular advancement for treatment of obstructive sleep apnea syndrome: a systematic review. *J Oral Maxillofac Surg* 69, e165–176 (2011).
- [67] Holty JE, Guilleminault C: Maxillomandibular advancement for the treatment of obstructive sleep apnea: a systematic review and meta-analysis. *Sleep Med Rev* 14, 287–297 (2010).
- [68] Vicini C, Dallan I, Campanini A, De Vito A, Barbanti F, Giorgiomarrano G et al.: Surgery vs ventilation in adult severe obstructive sleep apnea syndrome. *Am J Otolaryngol* 31, 14–20 (2010).

- [1] Blankenstein F: Wissenschaftliche Mitteilung der Deutschen Gesellschaft für zahnärztliche Prothetik und Werkstoffkunde (DGZPW) zur Verwendung thermoplastischer Nylon-Kunststoffe als Prothesenbasismaterial. *zm* 19, 42–44 (2009).
- [2] Blankenstein F: Alternative Prothesenbasiskunststoffe für Allergiker, Phobiker, Ästheten und Mikrostomiepatienten. *ZTM* 16 (3), 180–187 (2012).
- [3] Fuhr K, Behneke N, Reiber T: Die Teilprothese. München, Wien: Carl Hanser (1990).
- [4] Liebold Raff H, Wissing KH: Der Kommentar zu BEMA und GOZ. **Ort: Verlag** (2017).
- [5] Rzanny A, Göbel R, Nietzsche S, Küpper H: PEEK: Werkstoffkundliche Eigenschaften – mit Blick auf die dentale Anwendung. *ZTM* 21 (2), 102–110 (2017).
- [6] Schnettelker H: Die Geschichte der Kautschukprothese: Freiburg: Albert-Ludwigs-Universität (2001).
- [7] Wieckiewicz M, Opitz V, Richter G, Boening KW: Physical properties of polyamide-12 versus PMMA denture base materials. *BioMed Research International*, Article ID 150298 (2014).

- [1] Peppard PE, Young T, Barnet JH, Palta M, Hagen EW, Hla KM: Increased prevalence of sleep-disordered breathing in adults. *Am J Epidemiol* 177, 1006–1014 (2013).
- [2] Young T, Palta M, Dempsey J, Skatrud J, Weber S, Badr S: The occurrence of sleep-disordered breathing among middle-aged adults. *N Engl J Med* 328, 1230–1235 (1993).
- [3] Punjabi NM: The epidemiology of adult obstructive sleep apnea. *Proc Am Thorac Soc* 5, 136–143 (2008).
- [4] Young T, Peppard PE, Gottlieb DJ: Epidemiology of obstructive sleep apnea: a population health perspective. *Am J Respir Crit Care Med* 165, 1217–1239 (2002).
- [5] Sforza E, Roche F, Thomas-Anterion C, Kerleroux J, Beauchet O, Celle S et al.: Cognitive function and sleep related breathing disorders in a healthy elderly population: the SYNAPSE study. *Sleep* 33, 515–521 (2010).
- [6] Capistrano A, Cordeiro A, Capelozza Filho L, Almeida VC, Silva PI, Martinez S et al: Facial morphology and obstructive sleep apnea. *Dental Press J Orthod* 20, 60–67 (2015).
- [7] Denolf PL, Vanderveken OM, Marklund ME, Braem MJ: The status of cephalometry in the prediction of non-CPAP treatment outcome in obstructive sleep apnea patients. *Sleep Med Rev* 27, 56–73 (2016).
- [8] McNicholas WT: Diagnosis of obstructive sleep apnea in adults. *Proc Am Thorac Soc* 5, 154–160 (2008).
- [9] Ward Flemons W, McNicholas WT: Clinical prediction of the sleep apnea syndrome. *Sleep Med Rev* 1, 19–32 (1997).
- [10] Somers VK, White DP, Amin R, Abraham WT, Costa F, Culebras A et al.: Sleep apnea and cardiovascular disease: an American Heart Association/American College of Cardiology Foundation Scientific Statement from the American Heart Association Council for High Blood Pressure Research Professional Education Committee, Council on Clinical Cardiology, Stroke Council, and Council on Cardiovascular Nursing. *J Am Coll Cardiol* 52, 686–717 (2008).
- [11] Dong JY, Zhang YH, Qin LQ: Obstructive sleep apnea and cardiovascular risk: meta-analysis of prospective cohort studies. *Atherosclerosis* 229, 489–495 (2013).
- [12] Gami AS, Hodge DO, Herges RM, Olson EJ, Nykodym J, Kara T et al.: Obstructive sleep apnea, obesity, and the risk of incident atrial fibrillation. *J Am Coll Cardiol* 49, 565–571 (2007).
- [13] Gami AS, Pressman G, Caples SM, Kanagala R, Gard JJ, Davison DE et al.: Association of atrial fibrillation and obstructive sleep apnea. *Circulation* 110, 364–367 (2004).
- [14] Verse T, Dreher A, Heiser C, Herzog M, Maurer J, Pirsig W et al.: HNO-spezifische Therapie der obstruktiven Schlafapnoe bei Erwachsenen. AWMF-Register Nr. 017/069 (2015).
- [15] Geert M et al.: S3-Leitlinie Nicht erholsamer Schlaf/Schlafstörungen – Kapitel „Schlafbezogene Atmungsstörungen“. AWMF Registernummer 063/001 (2016).
- [16] Alves M, Jr., Franzotti ES, Baratieri C, Nunes LK, Nojima LI, Ruellas AC: Evaluation of pharyngeal airway space amongst different skeletal patterns. *Int J Oral Maxillofac Surg* 41, 814–819 (2012).
- [17] Abe-Nickler MD, Portner S, Sieg P, Hakim SG: No correlation between two-dimensional measurements and three-dimensional configuration of the pharyngeal upper airway space in cone-beam computed tomography. *J Craniomaxillofac Surg* 45, 371–376 (2017).
- [18] Abrahamyan L, Sahakyan Y, Chung S, Pechlivanoglou P, Bielecki J, Carcone SM et al.: Diagnostic accuracy of level IV portable sleep monitors versus polysomnography for obstructive sleep apnea: a systematic review and meta-analysis. *Sleep Breath* [AK1] (2018).
- [19] Danker-Hopfe H, Kunz D, Gruber G, Klosch G, Lorenzo JL, Himanen SL et al.: Interrater reliability between scorers from eight European sleep laboratories in subjects with different sleep disorders. *J Sleep Res* 13, 63–69 (2004).

- [20] Anandam A, Akinnusi M, Kufel T, Porhomayon J, El-Solh AA: Effects of dietary weight loss on obstructive sleep apnea: a meta-analysis. *Sleep Breath* 17, 227–234 (2013).
- [21] Benedict MA, Arterburn D: Worksite-based weight loss programs: a systematic review of recent literature. *Am J Health Promot* 22, 408–416 (2008).
- [22] Buchwald H, Avidor Y, Braunwald E, Jensen MD, Pories W, Fahrbach K et al.: Bariatric surgery: a systematic review and meta-analysis. *Jama* 292, 1724–1737 (2004).
- [23] Greenburg DL, Lettieri CJ, Eliasson AH: Effects of surgical weight loss on measures of obstructive sleep apnea: a meta-analysis. *Am J Med* 122, 535–542 (2009).
- [24] Sarkhosh K, Switzer NJ, El-Hadi M, Birch DW, Shi X, Karmali S: The impact of bariatric surgery on obstructive sleep apnea: a systematic review. *Obes Surg* 23, 414–423 (2013).
- [25] Permut I, Diaz-Abad M, Chatila W, Crocetti J, Gaughan JP, D'Alonzo GE et al.: Comparison of positional therapy to CPAP in patients with positional obstructive sleep apnea. *J Clin Sleep Med* 6, 238–243 (2010).
- [26] Skinner MA, Kingshott RN, Filsell S, Taylor DR: Efficacy of the 'tennis ball technique' versus nCPAP in the management of position-dependent obstructive sleep apnoea syndrome. *Respirology* 13, 708–715 (2008).
- [27] Ayas NT, FitzGerald JM, Fleetham JA, White DP, Schulzer M, Ryan CF et al.: Cost-effectiveness of continuous positive airway pressure therapy for moderate to severe obstructive sleep apnea/hypopnea. *Arch Intern Med* 166, 977–984 (2006).
- [28] Kushida CA, Littner MR, Hirshkowitz M, Morgenthaler TI, Alessi CA, Bailey D et al.: Practice parameters for the use of continuous and bilevel positive airway pressure devices to treat adult patients with sleep-related breathing disorders. *Sleep* 29, 375–380 (2006).
- [29] Randerath WJ, Hein H, Arzt M, Galetke W, Nilius G, Penzel T et al.: [Consensus paper on the diagnosis and treatment of sleep disordered breathing]. *Pneumologie* 68, 106–123 (2014).
- [30] Chai-Coetzer CL, Antic NA, Rowland LS, Reed RL, Esterman A, Catcheside PG et al.: Primary care vs specialist sleep center management of obstructive sleep apnea and daytime sleepiness and quality of life: a randomized trial. *Jama* 309, 997–1004 (2013).
- [31] Rosen CL, Auckley D, Benca R, Foldvary-Schaefer N, Iber C, Kapur V et al.: A multisite randomized trial of portable sleep studies and positive airway pressure autotitration versus laboratory-based polysomnography for the diagnosis and treatment of obstructive sleep apnea: the HomePAP study. *Sleep* 35, 757–767 (2012).
- [32] Gay P, Weaver T, Loubé D, Iber C: Evaluation of positive airway pressure treatment for sleep related breathing disorders in adults. *Sleep* 29, 381–401 (2006).
- [33] Giles TL, Lasserson TJ, Smith BH, White J, Wright J, Cates CJ: Continuous positive airways pressure for obstructive sleep apnoea in adults. *Cochrane Database Syst Rev*, Cd001106 (2006).
- [34] Sanders MH, Montserrat JM, Farre R, Givelber RJ: Positive pressure therapy: a perspective on evidence-based outcomes and methods of application. *Proc Am Thorac Soc* 5, 161–172 (2008).
- [35] Madbouly EM, Nadeem R, Nida M, Molnar J, Aggarwal S, Loomba R: The role of severity of obstructive sleep apnea measured by apnea-hypopnea index in predicting compliance with pressure therapy, a meta-analysis. *Am J Ther* 21, 260–264 (2014).
- [36] Wozniak DR, Lasserson TJ, Smith I: Educational, supportive and behavioural interventions to improve usage of continuous positive airway pressure machines in adults with obstructive sleep apnoea. *Cochrane Database Syst Rev*, Cd007736 (2014).
- [37] Kushida CA, Chediak A, Berry RB, Brown LK, Gozal D, Iber C et al.: Clinical guidelines for the manual titration of positive airway pressure in patients with obstructive sleep apnea. *J Clin Sleep Med* 4, 157–171 (2008).
- [38] Levy P, Kohler M, McNicholas WT, Barbe F, McEvoy RD, Somers VK et al.: Obstructive sleep apnoea syndrome. *Nat Rev Dis Primers* 1, 15015 (2015).
- [39] Weaver TE, Grunstein RR: Adherence to continuous positive airway pressure therapy: the challenge to effective treatment. *Proc Am Thorac Soc* 5, 173–178 (2008).

- [40] Epstein LJ, Kristo D, Strollo PJ, Jr., Friedman N, Malhotra A, Patil SP et al.: Clinical guideline for the evaluation, management and long-term care of obstructive sleep apnea in adults. *J Clin Sleep Med* 5, 263–276 (2009).
- [41] Li W, Xiao L, Hu J: The comparison of CPAP and oral appliances in treatment of patients with OSA: a systematic review and meta-analysis. *Respir Care* 58, 1184–1195 (2013).
- [42] Phillips CL, Grunstein RR, Darendeliler MA, Mihailidou AS, Srinivasan VK, Yee BJ et al.: Health outcomes of continuous positive airway pressure versus oral appliance treatment for obstructive sleep apnea: a randomized controlled trial. *Am J Respir Crit Care Med* 187, 879–887 (2013).
- [43] Randerath WJ, Verbraecken J, Andreas S, Bettge G, Boudewyns A, Hamans E et al.: Non-CPAP therapies in obstructive sleep apnoea. *Eur Respir J* 37, 1000–1028 (2011).
- [44] Ngiam J, Balasubramaniam R, Darendeliler MA, Cheng AT, Waters K, Sullivan CE: Clinical guidelines for oral appliance therapy in the treatment of snoring and obstructive sleep apnoea. *Aust Dent J* 58, 408–419 (2013).
- [45] Qaseem A, Holty JE, Owens DK, Dallas P, Starkey M, Shekelle P: Management of obstructive sleep apnea in adults: A clinical practice guideline from the American College of Physicians. *Ann Intern Med* 159, 471–483 (2013).
- [46] Puhan MA, Suarez A, Lo Cascio C, Zahn A, Heitz M, Braendli O: Didgeridoo playing as alternative treatment for obstructive sleep apnoea syndrome: randomised controlled trial. *Bmj* 332, 266–270 (2006).
- [47] Camacho M, Certal V, Abdullatif J, Zaghi S, Ruoff CM, Capasso R et al.: Myofunctional therapy to treat obstructive sleep apnea: a systematic review and meta-analysis. *Sleep* 38, 669–675 (2015).
- [48] Guimaraes KC, Drager LF, Genta PR, Marcondes BF, Lorenzi-Filho G: Effects of oropharyngeal exercises on patients with moderate obstructive sleep apnea syndrome. *Am J Respir Crit Care Med* 179, 962–966 (2009).
- [49] Randerath WJ, Galetke W, Domanski U, Weitkunat R, Ruhle KH: Tongue-muscle training by intraoral electrical neurostimulation in patients with obstructive sleep apnea. *Sleep* 27, 254–259 (2004).
- [50] Camacho M, Certal V, Brietzke SE, Holty JE, Guilleminault C, Capasso R: Tracheostomy as treatment for adult obstructive sleep apnea: a systematic review and meta-analysis. *Laryngoscope* 124, 803–811 (2014).
- [51] Browaldh N, Nerfeldt P, Lysdahl M, Bring J, Friberg D: SKUP3 randomised controlled trial: polysomnographic results after uvulopalatopharyngoplasty in selected patients with obstructive sleep apnoea. *Thorax* 68, 846–853 (2013).
- [52] Li HY, Wang PC, Lee LA, Chen NH, Fang TJ: Prediction of uvulopalatopharyngoplasty outcome: anatomy-based staging system versus severity-based staging system. *Sleep* 29, 1537–1541 (2006).
- [53] Marti S, Sampol G, Munoz X, Torres F, Roca A, Lloberes P et al.: Mortality in severe sleep apnoea/hypopnoea syndrome patients: impact of treatment. *Eur Respir J* 20, 1511–1518 (2002).
- [54] Goktas O, Solmaz M, Goktas G, Olze H: Long-term results in obstructive sleep apnea syndrome (OSAS) after laser-assisted uvulopalatoplasty (LAUP). *PLoS One* 9, e100211 (2014).
- [55] Sundaram S, Bridgman SA, Lim J, Lasserson TJ: Surgery for obstructive sleep apnoea. *Cochrane Database Syst Rev*, Cd001004 (2005).
- [56] Won CH, Li KK, Guilleminault C: Surgical treatment of obstructive sleep apnea: upper airway and maxillomandibular surgery. *Proc Am Thorac Soc* 5, 193–199 (2008).
- [57] Choi JH, Kim SN, Cho JH: Efficacy of the Pillar implant in the treatment of snoring and mild-to-moderate obstructive sleep apnea: a meta-analysis. *Laryngoscope* 123, 269–276 (2013).
- [58] Maurer JT, Sommer JU, Hein G, Hormann K, Heiser C, Stuck BA: Palatal implants in the treatment of obstructive sleep apnea: a randomised, placebo-controlled single-centre trial. *Eur Arch Otorhinolaryngol* 269, 1851–1856 (2012).

- [59] Back LJ, Hytonen ML, Roine RP, Malmivaara AO: Radiofrequency ablation treatment of soft palate for patients with snoring: a systematic review of effectiveness and adverse effects. *Laryngoscope* 119, 1241–1250 (2009).
- [60] Back LJ, Liukko T, Rantanen I, Peltola JS, Partinen M, Ylikoski J et al.: Radiofrequency surgery of the soft palate in the treatment of mild obstructive sleep apnea is not effective as a single-stage procedure: A randomized single-blinded placebo-controlled trial. *Laryngoscope* 119, 1621–1627 (2009).
- [61] Farrar J, Ryan J, Oliver E, Gillespie MB: Radiofrequency ablation for the treatment of obstructive sleep apnea: a meta-analysis. *Laryngoscope* 118, 1878–1883 (2008).
- [62] Strollo PJ, Jr., Soose RJ, Maurer JT, de Vries N, Cornelius J, Froymovich O et al.: Upper-airway stimulation for obstructive sleep apnea. *N Engl J Med* 370, 139–149 (2014).
- [63] Woodson BT, Gillespie MB, Soose RJ, Maurer JT, de Vries N, Steward DL et al.: Randomized controlled withdrawal study of upper airway stimulation on OSA: short- and long-term effect. *Otolaryngol Head Neck Surg* 151, 880–887 (2014).
- [64] Bangiyev JN, Traboulsi H, Abdulhamid I, Rozzelle A, Thottam PJ: Sleep architecture in Pierre-Robin sequence: The effect of mandibular distraction osteogenesis. *Int J Pediatr Otorhinolaryngol* 89, 72–75 (2016).
- [65] Vezina JP, Blumen M, Buchet I, Chabolle F: Sleep-disordered breathing: choosing the right cephalometric analysis. *J Oral Maxillofac Surg* 70, 1442–1448 (2012).
- [66] Pirklbauer K, Russmueller G, Stiebellehner L, Nell C, Sinko K, Millesi G et al.: Maxillomandibular advancement for treatment of obstructive sleep apnea syndrome: a systematic review. *J Oral Maxillofac Surg* 69, e165–176 (2011).
- [67] Holty JE, Guilleminault C: Maxillomandibular advancement for the treatment of obstructive sleep apnea: a systematic review and meta-analysis. *Sleep Med Rev* 14, 287–297 (2010).
- [68] Vicini C, Dallan I, Campanini A, De Vito A, Barbanti F, Giorgiomarrano G et al.: Surgery vs ventilation in adult severe obstructive sleep apnea syndrome. *Am J Otolaryngol* 31, 14–20 (2010).