

Sind Dentaleinheiten bereits vor der Installation mikrobiell kontaminiert?

- [1] Arbeitsgemeinschaft der Wissenschaftlichen Medizinischen Fachgesellschaften e.V.: Zahnärztliche Behandlungseinheiten, hygienische Anforderungen an das Wasser (2014); AWMF-Register-Nr. 075–002 (2014).
- [2] Costerton JW, Lewandowski Z, DeBeer D, Caldwell D, Korber D, James G: Biofilms, the customized microniche. *J of Bacteriology* 176, 2137–2142 (1994).
- [3] Sax H, Bloemberg G, Hasse B, Sommerstein R, Kohler P, Achermann Y, Rössle M, Falk V, Kuster SP, Böttger EC, Weber R: Prolonged outbreak of *Mycobacterium chimaera* infection after open-chest heart surgery. *Clin Infect Dis* 61 (1), 67–75 (2015).
- [4] Sommerstein R, Rüegg C, Kohler P, Bloemberg G, Kuster SP, Sax H: Transmission of *Mycobacterium chimaera* from heater-cooler units during cardiac surgery despite an ultraclean air ventilation system. *Emerg Infect* 22 (6), 1008–1013 (2016).
- [5] Sorin Group Deutschland GmbH: Sicherheitshinweis zur Desinfektion und Reinigung von Hypothermiegeräten 1T und 3T. 08.06.2015. Referenznummer 1223/15.
- [6] Lück C, Neumann K, Reitemeier B, Jatzwauk L: Legionellen in Dentaleinheiten – Infektionsrisiko für Patienten und Personal? *Zahnmedizin up2date* 4, 345–359 (2011).

Direkte adhäsive Versorgungen wurzelkanalbehandelter Zähne

- [1] Da Rosa Rodolfo PA, Donassollo TA, Cenci MS, Loguércio AD, Moraes RR, Bronkhorst EM, Opdam NJ, Demarco FF. 22 year clinical evaluation of the performance of two posterior composites with different filler characteristics. Dent Mater 2011; 27: 955–963.
- [2] Dammaschke T, Nykiel K, Sagheri D, Schäfer E. Influence of coronal restorations on the fracture resistance of root canal-treated premolar and molar teeth: a retrospective study. Aust Endod J 2013; 39: 48–56.
- [3] Deliperi S, Bardwell DN. Direct cuspal-coverage posterior resin composite restorations: A case report. Oper Dent 2006; 31: 143–150.
- [4] ElAyouti A, Serry MI, Geis-Gerstorfer J, Löst C. Influence of cusp coverage on the fracture resistance of premolars with endodontic access cavities. Int Endod J 2011; 44: 543–549.
- [5] Ernst CP. Die 20 beliebtesten Fehler beim Kleben. DFZ 2010; 11: 66–83.
- [6] Ernst CP. CEREC versus Straumann: Welche Teilkronen passt besser? Teil 1. ZMK 2013; 29: 184–193.
- [7] Ernst CP. Die Verwendung eines weiß-opaken Flow-Kompositen zur Markierung tiefer Kavitätenanteile in der Klasse II. Ästhetische Zahnmedizin 2004; 7: 160–169.
- [8] Ernst CP, Rullmann I, Janssen B, Willershausen B. Polymerization shrinkage stress of bulk fill resin composites. Vortrag auf der IADR Jahrestagung 2014. <https://iadr.confex.com/iadr/13iags/webprogram/Paper170911.html>.
- [9] Ferracane J, Watts DC, Barghi N, Ernst CP, Rueggeberg FA, Shortall A, Price R, Strassler H. Der effiziente Einsatz von Lichtpolymerisationsgeräten – ein Leitfaden für Zahnärzte. ZMK 2014; 30: 166–180.
- [10] Ferrari M, Cagidiaco MC, Goracci C, Vichi A, Mason PN, Radovic I, Tay F. Long-term retrospective study of the clinical performance of fiber posts. Am J Dent 2007; 20:287–291.
- [11] Frankenberger R, Reinelt C, Roggendorf MJ, Lrämer N. Direkte oder indirekte Restaurationen im Seitenzahnbereich? Quintessenz 2010; 561: 575–579.
- [12] Goracci C, Grandini S, Bossù M, Bertelli E, Ferrari M. Laboratory assessment of the retentive potential of adhesive posts: a review. J Dent 2007; 35: 827–835.
- [13] Hickel R, Ernst CP, Haller B, Hugo B, Kunzelmann KH, Merte K, Ott K, Schmalz G, Staehle HJ. Direkte Kompositrestaurationen im Seitenzahnbereich – Indikation und Lebensdauer Zahnärztliche Mitteilungen 2005; 95: 74–76.
- [14] Juloski J, Fadda GM, Monticelli F, Fajó-Pascual M, Goracci C, Ferrari M. Four-year Survival of Endodontically Treated Premolars Restored with Fiber Posts. J Dent Res 2014 19; 93 (7 suppl): 52S–58S.
- [15] Kon M, Nicola Zitzmann NU, Weiger R, Krastl G. Postendodontische Restauration: eine Umfrage unter Zahnärzten der Schweiz. Schweiz Monatsschr Zahnmed 2013; 123: 1083–1088.
- [16] Manhart, J, Chen H, Hamm, G, Hickel R: Buonocore Memorial Lecture. Review of the clinical survival of direct and indirect restorations in posterior teeth of the permanent dentition. Oper Dent 2004;29: 481–508.
- [17] Mannocci F, Bertelli E, Sherriff M, Watson TF, Ford TR. Three-year clinical comparison of survival of endodontically treated teeth restored with either full cast coverage or with direct composite restoration. J Prosthet Dent 2002; 88:297–301.
- [18] Mohammadi N, Kahnamoii MA, Yeganeh PK, Navimipour EJ. Effect of fiber post and cusp coverage on fracture resistance of endodontically treated maxillary premolars directly restored with composite resin. J Endod. 2009; 35: 1428–1432.

- [19] [Mondelli RF](#), [Ishikirama SK](#), [de Oliveira Filho O](#), [Mondelli J](#). Fracture resistance of weakened teeth restored with condensable resin with and without cusp coverage. *J Appl Oral Sci* 2009;17: 161–165.
- [20] Nagasiri R, Chitmongkolsuk S: Long-term survival of endodontically treated molars without crown coverage: a retrospective cohort study. *J Prosthet Dent.* 2005; 93: 164–170.
- [21] [Naumann M](#), [Preuss A](#), [Rosentritt M](#). Effect of incomplete crown ferrules on load capacity of endodontically treated maxillary incisors restored with fiber posts, composite build-ups, and all-ceramic crowns: an in vitro evaluation after chewing simulation. *Acta Odontol Scand* 2006; 64: 31–36.
- [22] [Naumann M](#), [Koelpin M](#), [Beuer F](#), [Meyer-Lueckel H](#). 10-year survival evaluation for glass-fiber-supported postendodontic restoration: a prospective observational clinical study. *J Endod* 2012; 38: 432–435.
- [23] [Nothdurft FP](#), [Seidel E](#), [Gebhart F](#), [Naumann M](#), [Motter PJ](#), [Pospiech PR](#). Influence of endodontic posts on the fracture behavior of crowned premolars with Class II cavities. *J Dent* 2008; 36: 287–393.
- [24] Opdam NJM, Bronkhorst EM, Loomans BAC, Huysmans M-CDNJM: 12-year survival of composite vs. amalgam restorations. *JDR Online First*, published on July 26, 2010.
- [25] Plotino G, Buono L, Grande NM, Lsamorgese V, Somma F. Fracture resistance of endodontically treated molars restored with extensive composite resin restorations. *J Prosthet Dent* 2008; 99: 225–232.
- [26] Rosentritt M, Behr M, Kolbeck C, Handel G. Flexural strength of restorative composites after different aging conditions. Vortrag auf der IADR Jahrestagung 2014. <https://iadr.confex.com/iadr/13iags/webprogram/Paper171330.html>
- [27] Rullmann I, Schattenberg A, Marx M, Willershausen B, Ernst CP: Spannungsoptische Messungen der Polymerisationsschrumpfungskraft schrumpfreduzierter Komposite. *Schweiz Monatsschr Zahnmed* 2012; 122: 8–12.
- [28] Scotti N, Rota R, Scanetti M, Paolono DS, Chisandussi G, Pasqualini D, Berutti E. Influence of adhesive techniques on fracture resistance of endodontically treated premolars with various residual wall thicknesses. *J Prosthet Dent* 2013; 110: 376382.
- [29] Staehle J. Restauration eines “hoffnungslosen Zahnes” mittels direkter Kompositkrone. *Quintessenz* 2006; 57: 531–537.
- [30] [Sterzenbach G](#), [Franke A](#), [Naumann M](#). Rigid versus flexible dentine-like endodontic posts-clinical testing of a biomechanical concept: seven-year results of a randomized controlled clinical pilot trial on endodontically treated abutment teeth with severe hard tissue loss. *J Endod* 2012; 38: 1557–1563.
- [31] [Xie KX](#), [Wang XY](#), [Gao XJ](#), [Yuan CY](#), [Li JX](#), [Chu CH](#). Fracture resistance of root filled premolar teeth restored with direct composite resin with or without cusp coverage. *Int Endod J.* 2012; 45: 524–529.

- [1] Bender W, Taubenheim L: Intraligamentäre Anästhesie – Eine ernst zu nehmende Alternative? Zahnärztl Mitt 4, 48–52 (2016).
- [2] Benz C, Prothmann M, Taubenheim L: Die intraligamentäre Anästhesie – Primäre Methode der dentalen Lokalanästhesie. Deutscher Zahnärzte Verlag, Köln (2015).
- [3] BGB § 630 – Patientenrechtegesetz (2013).
- [4] Csides M, Taubenheim L, Glockmann E: Intraligamentäre Anästhesie: Grenzen und Komplikationen. Dtsch Zahnärztl Z 8, 561–569 (2011).
- [5] Eifinger FF, Fritz U, Gautzsch F: Untersuchung zur passiven Aspiration bei der Lokalanästhesie. ZWR Deutsch Zahnärztlebl 3, 166–1176 (1983).
- [6] Gray RJM, Lomax AM, Rood JP: Periodontal ligament injection: with or without a vasoconstrictor? Br Dent J 162, 263–265 (1987).
- [7] Lipp MDW: Lokalanästhesie – ohne Risiko für jeden Patienten? Zahnärztl Prax 9, 330–333 (1989).
- [8] Prothmann M, Taubenheim L, Benz C: Die intraligamentäre Anästhesie in der zahnärztlichen Praxis – Metaanalyse der klinischen Erfahrungen von 1976 bis 2014. Deutsch Zahnärztl Z 71 (2), 136–145 (2016).
- [9] Rahn R: Techniken der Lokalanästhesie. forum_med_dent; Aventis, 82–101 (2001).
- [10] Schwenzer N, Ehrenfeld M: Zahn-Mund-Kiefer-Heilkunde, Bd. 3: Zahnärztl Chirurgie. Thieme, 2000, Stuttgart, New York.
- [11] Taubenheim L, Bayer G: Intraligamentäre Anästhesie – primäre Methode der zahnärztlichen örtlichen Betäubung. i-med Akademie FORUM Verlag 2014; ISBN 978 3 86586 479 6.
- [12] Zugal W, Taubenheim L, Schulz D: Triade des Anästhesie-Erfolgs: Instrumente – Anästhetika – Methoden Beherrschung. Z Stomatol 102.1, 9–14 (2005).

Entscheidungsparameter für die Ex- oder NonEx-Therapie in der Kieferorthopädie

- [1] Angle EH: Treatment of malocclusions of teeth. 7th ed. S.S. White Dental Manufacturing Comp, Philadelphia (1907).
- [2] Begg PR: Stone age man's dentition. Am J Orthod 40, 298–312, 373–383, 462–475, 517–531 (1954).
- [3] Berseth A: The stability of the lower dental arch orthodontic treatment. Europ Orthodont Soc Trans 251 (1972).
- [4] Björk A: Prediction of mandibular growth rotation. Am J Orthod 55, 585 (1969).
- [5] Bolton WA: Disharmony in tooth size and its relation to the analysis and treatment of malocclusion. Angle Orthod 28, 113–130 (1958).
- [6] Corruccini RS: Australian aboriginal tooth succession, interproximal attrition and Begg's theory. Am J Orthod Dentofacial Orthop 97 (4), 349–57 (1990).
- [7] Derichsweiler H: Die Gaumennahterweiterung. Fortschr Kieferorthop 14, 5 (1953).
- [8] Franchi L, Baccetti T, McNamara J: Cephalometric floating norms for North American adults. Angle Orthod 68 (6), 497–502 (1998).
- [9] Gross A, Hasund A: Neuere vergleichende korrelationsstatistische Untersuchungen zur Vorhersage des Platzbedarfs in den Stützonen durch multiple Regressionsgleichungen. Fortschr Kieferorthop 50, 109 (1989).
- [10] Habersack K, Hasund A: Clinical application of individualized cephalometrics. CEPH TEMPLATE a practical tool. SAM Präzisionstechnik GmbH, Gauting (2013).
- [11] Habersack K, Karoglan A: Stützonenprognose im Unterkiefer. Treffsicherheit von sechs Analysen im Vergleich. Poster DGKFO München (2003).
- [12] Habersack K, Karoglan A, Sommer B, Benner KU: High-resolution multislice computerized tomography with multiplanar and 3-dimensional reformation imaging in rapid palatal expansion. Am J Orthod Dentofacial Orthop 131, 776–781, (2007).
- [13] Habersack K, Ramian C: Die HANSA IV-Schiene. Quintessenz Zahntech 39 (11), 1652–1660 (2013).
- [14] Hasund A: Okklusjon og facalkranium i middelalderbefolkningen i Oslo og Heidal: en odontometrisk, kraniometrisk og røntgenkefalometrisk studie. Habilchrift Oslo (1966).
- [15] Hasund A: Attrition and dental arch space. Trans Eur Orthod Soc, 131–142 (1965).
- [16] Hasund A: Klinische Kephalometrie für die Bergen-Technik. Kieferorthopädische Abteilung des zahnärztlichen Instituts der Universität Bergen, Norwegen (1974).
- [17] Hasund A, Bingler P, Rudzki I: Edgewise-Technik. Standard-Edgewise und Straight-wire-System. Praxis der Zahnheilkunde, Kieferorthopädie II, Hrsg. P. Diedrich Urban & Fischer (Studiennachdruck) (2010).
- [18] Hasund A, Böe OE: Floating norms as a guidance for the position of the lower incisor. Angle Orthod 50, 165–168 (1980).
- [19] Hasund A, Böe OE, Wisth PJ: Der H-Winkel in der kieferorthopädischen Diagnostik. Fortschr Kieferorthop. 41, 40–46 (1980).
- [20] Hasund A, Ulstein G: The position of the incisors in relation to the lines NA and NB in different facial types. Am J Orthod 57, 1–14 (1970).
- [21] Hotz R: Guidance of eruption versus serial extraction. Am J Orthod 58, 1–20 (1970).
- [22] Kjellgren B: Serial extraction as a corrective procedure in dental orthopedic therapy. Acta Odontol Scand 8, 17–43 (1948).
- [23] Krogh-Poulsen W, Olsson A: Occlusal disharmonies and dysfunction of the stomatognathic system. Dental Clinics of North America 10, 627 (1966).
- [24] Riedel RA, Little RM, Bui TD: Mandibular incisor extraction – postretention evaluation of stability and relapse. Angle Orthod 62 (2), 103–16 (1992).
- [25] Segner D, Hasund A: Individualisierte Kephalometrie. Segner, HansaDont Verlag und Vertrieb, Hamburg (1991).

- [26] Solow B: Die dentoalveolare Kompensation: Grundlagen und klinische Bedeutung. *Inf Orthod Kieferorthop* 38, 148–162 (2006).
- [27] Steiner C: Cephalometrics for you and me. *Am J Orthod* 39, 729–755 (1953).
- [28] Steinhäuser E, Janson I: Kieferorthopädische Chirurgie. Eine interdisziplinäre Aufgabe. Band I. Quintessenz Verlag Berlin (1988).
- [29] Tollaro I, Baccetti T, Franchi L: Floating norms for the assessment of craniofacial pattern in the deciduous dentition. *Eur J Orthod* 18, 359–365 (1996).
- [30] Weise W: Kieferorthopädische Kombinationstherapie. München, Wien, Baltimore: Urban und Schwarzenberg (1992).

Low-Level-Laser-Therapie/Laser-Photo-Therapie in der Zahnheilkunde – eine Übersicht

- [1] Mester E: The effect of laser beams on the grow of hair in mice. Radiobiol Radiother 9, 621–626 (1968).
- [2] Mester E: Über die Wirkungen von Laserstrahlen auf die Bakterienphagozytose der Leukozyten. Acta biol Med germ 21, 317–324 (1968).
- [3] Mester E: Clinical application of Laser beams. Lyon Chir 65, 335–345 (1969).
- [4] Mester E: Effect of laser rays on muscle fiber regeneration. Acta Chir Acad Sci Hung 13, 315–324 (1974).
- [5] Mester E: Laser-induce stimulation of the vascularisation of healing wound. Separat Exp 30, 341–345 (1974).
- [6] Gaus H: Der Low-Level-Laser in der Zahnarztpraxis. Laser Journal 1, 22ff; Laser Journal 2, 27ff (2007).
- [7] Warnke U: Der Diodenlaser. Deutsches Ärzteblatt 84 (44), 29ff (1987).
- [8] Roynesdal AK et al: The effect of soft laser application on postoperative pain and swelling. A double-blind, crossover study. Int J Oral Maxillofac Surg 22, 242–245 (1993).
- [9] Ferrante M et al: Effect of low-level-laser-therapy after extraction of impacted lower third molars. Lasers in Medical Science 28 (3), 845–849 (2013).
- [10] Neckel Cl: Biostimulation: A comparative study in the postoperative outcome of patients after third molar extraction. Journal of Oral Laser Applications 1 (3), 215–219 (2001).
- [11] Fernando S: A randomised double blind comparative study of low level laser therapy following surgical extraction of lower third molar teeth. Br J Oral Maxillofac Surg 31, 170–172 (1993).
- [12] Taube S: Helium-neon laser therapy in the prevention of postoperative swelling and pain after wisdom tooth extraction. Proc Finn Dent 86, 23–27 (1990).
- [13] Bramante CM et al: Use of a 660 nm Laser to Aid in the Healing of Necrotic Alveolar Mucosa caused by Extruded Sodium Hypochlorite: A Case Report. J Endod 41 (11), 1899–1902 (2015).
- [14] Munoz PJ et al: The Effect of 670 nm Low Laser Therapy on Herpes Simplex Type 1. Photomed Laser Surg 30 (1), 37–40 (2012).
- [15] Schindl A, Neumann R: Low-Intensity Laser Therapy Is An Effective Treatment For Recurrent Herpes Simplex Infection. Investigative Dermatology 113 (2), 221–223 (1999).
- [16] Chow Roberta T et al: Efficacy of low-level-laser-therapy in the management of neck-pain: a systematic review and meta-analysis of randomised placebo or active treatment controlled trials. The Lancet 374 (9705), 1897–1908 (2009).
- [17] Sayed N et al: Management of Temporomandibular Disorders with Low Level Laser Therapy. J Maxillofac Oral Surg 13, 444–450 (2014).
- [18] Gungor A: Effectiveness of transmeatal low power laser irradiation for chronic tinnitus. J Laryngol Otol 122 (5), 447–451 (2008).
- [19] Cuda D, De Caria A: Effectiveness of combined counselling and low level laser stimulation in the treatment of disturbing chronic tinnitus. Int Tinnitus Journal 2, 175–180 (2008).