

- [1] Anderson LE, Inglehart MR, El-Kholy K, Eber R, Wang HL: Implant associated soft tissue defects in anterior maxilla: a randomized control trial comparing subepithelial connective tissue graft and acellular dermal matrix allograft.: *Implant Dent* (4),416–25 (2014).
- [2] Araujo MG, Lindhe J: Dimensional ridge alterations following tooth extraction. An experimental study in the dog. *J Clin Periodontol* 32, 212–218 (2005).
- [3] Araujo M, Linder E, Wennström J, Lindhe J: The Influence of Bio-Oss Collagen on Healing of an Extraction Socket: An Experimental Study in the Dog. *Int J Periodontics Restorative Dent* 28,123–135 (2008).
- [4] Basetti RG, Stähli A, Basetti MA, Scuean A: Soft tissue augmentation procedures at second- stage surgery: a systematic review. *Clin Oral Investig* 2016 Apr. 4. (Epub ahead of print).
- [5] Block MS et al.: Hydroxyapatite-coated cylindrical implants in the posterior mandible: 10-year observations. *Int J Oral Maxillofac Implants* 11, 626–633 (1996).
- [6] Cardaropoli D, Cardaropoli G: Preservation of the postextraction alveolar ridge: a clinical and histologic study. *Int J Periodontics Restorative Dent* 28 (5), 469–477 (2008).
- [7] Chiu YW, Lee SY, Lin YC, Lai YL: Significance of the width of keratinized mucosa on peri-implant health. *J Chin Med Assoc* 78 (7), 389–94 (2015).
- [8] Deporter DA et al: A histological evaluation of a functional endosseous, porous-surfaced, titanium alloy dental implant system in the dog. *J Dent Res* 67, 1190–1195 (1988).
- [9] Fickl S: Peri-implant mucosal recession: Clinical significance and therapeutic opportunities. *Quintessence Int* 46 (8), 671–6 (2015).
- [10] Kirsch A, Ackermann KL: The IMZ osteointegrated implant system. *Dent Clin North Am* 33, 733–791 (1989).
- [11] Lee JH, Lee JS, Baek WS, Lim HC, Cha JK, Choi SH, Jung UW: Assessment of dehydrothermally cross-linked collagen membrane for guided bone regeneration around peri-implant dehiscence defects: a randomized single-blinded clinical trial. *J Periodontal Implant Sci* 45 (6), 229–37 (2015).
- [12] Lima RS, Peruzzo DC, Napimoga MH, Saba-Chuifi E, Santos-Pereira SA, Martinez EF: Evaluation of the Biological Behavior of Mucograft in Human Gingival Fibroblasts: An In Vitro Study. *Braz Dent J* 26 (6), 602–6 (2015).
- [13] Lin GH, Chan HL, Wang HLJ: The significance of keratinized mucosa on implant health: a systematic review. *J Periodontol* 84 (12), 1755–67 (2013).
- [14] Liu C, Su Y, Tan B, Ma P, Wu G, Li J, Geng W: Reconstruction of attached soft tissue around dental implants by acellular dermal matrix grafts and resin splint. *Int J Clin Exp Med* 7 (12), 4666–76 (2014).
- [15] McGuire MK, Scheyer ET: Xenogeneic collagen matrix with coronally advanced flap compared to connective tissue with coronally advanced flap for the treatment of dehiscence-type recession defects. *J Periodontol* 81 (8), 1108–17 (2010).
- [16] Roos-Jansaker AM et al: Nine-to fourteen-year follow-up of implant treatment. Part I: implant loss and associations to various factors. *J Clin Periodontol* 33, 283–289 (2006).
- [17] Roos-Jansaker AM et al: Nine-to fourteen-year follow-up of implant treatment. Part II: presence of peri-implant lesions. *J Clin Periodontol* 33, 290–295 (2006).
- [18] Roos-Jansaker AM et al: Nine-to fourteen-year follow-up of implant treatment. Part I: factors associated with peri-implant lesions. *J Clin Periodontol* 33, 296–301 (2006).

- [19] Sanz M et al: Clinical evaluation of a new collagen matrix (Mucograft prototype) to enhance the width of keratinized tissue in patients with fixed prosthetic restorations: a randomized prospective clinical trial. *J Clin Periodontol* 36 (10), 868–76 (2009).
- [20] Schmitt CM, Moest T, Lutz R, Werhahn F, Neukam FW, Schlegel KA: Long-term outcomes after vestibuloplasty with a porcine collagen matrix (Mucograft) versus the free gingival graft: a comparative prospective clinical trial. *Clin Oral Implants res.* 2015 Feb 27. (Epub ahead of print).
- [21] Schropp L, Wenzel A, Kostopoulos L, Karring T: Bone healing soft tissue contour changes followig single-tooth extraction: A clinical and radiographic 12-month prospective study. *Int J Periodontics Restorative Dent* 23, 313–323 (2003).
- [22] Strietzel FP, Shakibaie MB: Der Einsatz der TefGen-FD[®]-Membran zum Erhalt des Alveolarkamms nach Zahnextraktionen: eine klinische Studie. *Dtsch Zahnärztl Z* 12, 883–886 (1998).