

**Therapieoptionen der obstruktiven Schlapnoe**

- [1] Peppard PE, Young T, Barnet JH, Palta M, Hagen EW, Hla KM: Increased prevalence of sleep-disordered breathing in adults. *Am J Epidemiol* 177, 1006–1014 (2013).
- [2] Young T, Palta M, Dempsey J, Skatrud J, Weber S, Badr S: The occurrence of sleep-disordered breathing among middle-aged adults. *N Engl J Med* 328, 1230–1235 (1993).
- [3] Punjabi NM: The epidemiology of adult obstructive sleep apnea. *Proc Am Thorac Soc* 5, 136–143 (2008).
- [4] Young T, Peppard PE, Gottlieb DJ: Epidemiology of obstructive sleep apnea: a population health perspective. *Am J Respir Crit Care Med* 165, 1217–1239 (2002).
- [5] Sforza E, Roche F, Thomas-Anterion C, Kerleroux J, Beauchet O, Celle S et al.: Cognitive function and sleep related breathing disorders in a healthy elderly population: the SYNAPSE study. *Sleep* 33, 515–521 (2010).
- [6] Capistrano A, Cordeiro A, Capelozza Filho L, Almeida VC, Silva PI, Martinez S et al: Facial morphology and obstructive sleep apnea. *Dental Press J Orthod* 20, 60–67 (2015).
- [7] Denolf PL, Vanderveken OM, Marklund ME, Braem MJ: The status of cephalometry in the prediction of non-CPAP treatment outcome in obstructive sleep apnea patients. *Sleep Med Rev* 27, 56–73 (2016).
- [8] McNicholas WT: Diagnosis of obstructive sleep apnea in adults. *Proc Am Thorac Soc* 5, 154–160 (2008).
- [9] Ward Flemons W, McNicholas WT: Clinical prediction of the sleep apnea syndrome. *Sleep Med Rev* 1, 19–32 (1997).
- [10] Somers VK, White DP, Amin R, Abraham WT, Costa F, Culebras A et al.: Sleep apnea and cardiovascular disease: an American Heart Association/American College of Cardiology Foundation Scientific Statement from the American Heart Association Council for High Blood Pressure Research Professional Education Committee, Council on Clinical Cardiology, Stroke Council, and Council on Cardiovascular Nursing. *J Am Coll Cardiol* 52, 686–717 (2008).
- [11] Dong JY, Zhang YH, Qin LQ: Obstructive sleep apnea and cardiovascular risk: meta-analysis of prospective cohort studies. *Atherosclerosis* 229, 489–495 (2013).
- [12] Gami AS, Hodge DO, Herges RM, Olson EJ, Nykodym J, Kara T et al.: Obstructive sleep apnea, obesity, and the risk of incident atrial fibrillation. *J Am Coll Cardiol* 49, 565–571 (2007).
- [13] Gami AS, Pressman G, Caples SM, Kanagala R, Gard JJ, Davison DE et al.: Association of atrial fibrillation and obstructive sleep apnea. *Circulation* 110, 364–367 (2004).
- [14] Verse T, Dreher A, Heiser C, Herzog M, Maurer J, Pirsig W et al.: HNO-spezifische Therapie der obstruktiven Schlapnoe bei Erwachsenen. *AWMF-Register Nr. 017/069* (2015).
- [15] Geert M et al.: S3-Leitlinie Nicht erholsamer Schlaf/Schlafstörungen – Kapitel „Schlafbezogene Atmungsstörungen“. *AWMF Registernummer 063/001* (2016).
- [16] Alves M, Jr., Franzotti ES, Baratieri C, Nunes LK, Nojima LI, Ruellas AC: Evaluation of pharyngeal airway space amongst different skeletal patterns. *Int J Oral Maxillofac Surg* 41, 814–819 (2012).
- [17] Abe-Nickler MD, Portner S, Sieg P, Hakim SG: No correlation between two-dimensional measurements and three-dimensional configuration of the pharyngeal upper airway space in cone-beam computed tomography. *J Craniomaxillofac Surg* 45, 371–376 (2017).
- [18] Abrahamyan L, Sahakyan Y, Chung S, Pechlivanoglou P, Bielecki J, Carcone SM et al.: Diagnostic accuracy of level IV portable sleep monitors versus polysomnography for obstructive sleep apnea: a systematic review and meta-analysis. *Sleep Breath* [AK1] (2018).
- [19] Danker-Hopfe H, Kunz D, Gruber G, Klosch G, Lorenzo JL, Himanen SL et al.: Interrater reliability between scorers from eight European sleep laboratories in subjects with different sleep disorders. *J Sleep Res* 13, 63–69 (2004).

- [20] Anandam A, Akinnusi M, Kufel T, Porhomayon J, El-Soh AA: Effects of dietary weight loss on obstructive sleep apnea: a meta-analysis. *Sleep Breath* 17, 227–234 (2013).
- [21] Benedict MA, Arterburn D: Worksite-based weight loss programs: a systematic review of recent literature. *Am J Health Promot* 22, 408–416 (2008).
- [22] Buchwald H, Avidor Y, Braunwald E, Jensen MD, Pories W, Fahrbach K et al.: Bariatric surgery: a systematic review and meta-analysis. *Jama* 292, 1724–1737 (2004).
- [23] Greenburg DL, Lettieri CJ, Eliasson AH: Effects of surgical weight loss on measures of obstructive sleep apnea: a meta-analysis. *Am J Med* 122, 535–542 (2009).
- [24] Sarkhosh K, Switzer NJ, El-Hadi M, Birch DW, Shi X, Karmali S: The impact of bariatric surgery on obstructive sleep apnea: a systematic review. *Obes Surg* 23, 414–423 (2013).
- [25] Permut I, Diaz-Abad M, Chatila W, Crocetti J, Gaughan JP, D'Alonzo GE et al.: Comparison of positional therapy to CPAP in patients with positional obstructive sleep apnea. *J Clin Sleep Med* 6, 238–243 (2010).
- [26] Skinner MA, Kingshott RN, Filsell S, Taylor DR: Efficacy of the 'tennis ball technique' versus nCPAP in the management of position-dependent obstructive sleep apnoea syndrome. *Respirology* 13, 708–715 (2008).
- [27] Ayas NT, FitzGerald JM, Fleetham JA, White DP, Schulzer M, Ryan CF et al.: Cost-effectiveness of continuous positive airway pressure therapy for moderate to severe obstructive sleep apnea/hypopnea. *Arch Intern Med* 166, 977–984 (2006).
- [28] Kushida CA, Littner MR, Hirshkowitz M, Morgenthaler TI, Alessi CA, Bailey D et al.: Practice parameters for the use of continuous and bilevel positive airway pressure devices to treat adult patients with sleep-related breathing disorders. *Sleep* 29, 375–380 (2006).
- [29] Randerath WJ, Hein H, Arzt M, Galetke W, Nilius G, Penzel T et al.: [Consensus paper on the diagnosis and treatment of sleep disordered breathing]. *Pneumologie* 68, 106–123 (2014).
- [30] Chai-Coetzer CL, Antic NA, Rowland LS, Reed RL, Esterman A, Catcheside PG et al.: Primary care vs specialist sleep center management of obstructive sleep apnea and daytime sleepiness and quality of life: a randomized trial. *Jama* 309, 997–1004 (2013).
- [31] Rosen CL, Auckley D, Benca R, Foldvary-Schaefer N, Iber C, Kapur V et al.: A multisite randomized trial of portable sleep studies and positive airway pressure autotitration versus laboratory-based polysomnography for the diagnosis and treatment of obstructive sleep apnea: the HomePAP study. *Sleep* 35, 757–767 (2012).
- [32] Gay P, Weaver T, Loube D, Iber C: Evaluation of positive airway pressure treatment for sleep related breathing disorders in adults. *Sleep* 29, 381–401 (2006).
- [33] Giles TL, Lasserson TJ, Smith BH, White J, Wright J, Cates CJ: Continuous positive airways pressure for obstructive sleep apnoea in adults. *Cochrane Database Syst Rev*, Cd001106 (2006).
- [34] Sanders MH, Montserrat JM, Farre R, Givelber RJ: Positive pressure therapy: a perspective on evidence-based outcomes and methods of application. *Proc Am Thorac Soc* 5, 161–172 (2008).
- [35] Madbouly EM, Nadeem R, Nida M, Molnar J, Aggarwal S, Loomba R: The role of severity of obstructive sleep apnea measured by apnea-hypopnea index in predicting compliance with pressure therapy, a meta-analysis. *Am J Ther* 21, 260–264 (2014).
- [36] Wozniak DR, Lasserson TJ, Smith I: Educational, supportive and behavioural interventions to improve usage of continuous positive airway pressure machines in adults with obstructive sleep apnoea. *Cochrane Database Syst Rev*, Cd007736 (2014).
- [37] Kushida CA, Chediak A, Berry RB, Brown LK, Gozal D, Iber C et al.: Clinical guidelines for the manual titration of positive airway pressure in patients with obstructive sleep apnea. *J Clin Sleep Med* 4, 157–171 (2008).
- [38] Levy P, Kohler M, McNicholas WT, Barbe F, McEvoy RD, Somers VK et al.: Obstructive sleep apnoea syndrome. *Nat Rev Dis Primers* 1, 15015 (2015).
- [39] Weaver TE, Grunstein RR: Adherence to continuous positive airway pressure therapy: the challenge to effective treatment. *Proc Am Thorac Soc* 5, 173–178 (2008).

- [40] Epstein LJ, Kristo D, Strollo PJ, Jr., Friedman N, Malhotra A, Patil SP et al.: Clinical guideline for the evaluation, management and long-term care of obstructive sleep apnea in adults. *J Clin Sleep Med* 5, 263–276 (2009).
- [41] Li W, Xiao L, Hu J: The comparison of CPAP and oral appliances in treatment of patients with OSA: a systematic review and meta-analysis. *Respir Care* 58, 1184–1195 (2013).
- [42] Phillips CL, Grunstein RR, Darendeliler MA, Mihailidou AS, Srinivasan VK, Yee BJ et al.: Health outcomes of continuous positive airway pressure versus oral appliance treatment for obstructive sleep apnea: a randomized controlled trial. *Am J Respir Crit Care Med* 187, 879–887 (2013).
- [43] Randerath WJ, Verbraecken J, Andreas S, Bettega G, Boudewyns A, Hamans E et al.: Non-CPAP therapies in obstructive sleep apnoea. *Eur Respir J* 37, 1000–1028 (2011).
- [44] Ngiam J, Balasubramaniam R, Darendeliler MA, Cheng AT, Waters K, Sullivan CE: Clinical guidelines for oral appliance therapy in the treatment of snoring and obstructive sleep apnoea. *Aust Dent J* 58, 408–419 (2013).
- [45] Qaseem A, Holty JE, Owens DK, Dallas P, Starkey M, Shekelle P: Management of obstructive sleep apnea in adults: A clinical practice guideline from the American College of Physicians. *Ann Intern Med* 159, 471–483 (2013).
- [46] Puhan MA, Suarez A, Lo Cascio C, Zahn A, Heitz M, Braendli O: Didgeridoo playing as alternative treatment for obstructive sleep apnoea syndrome: randomised controlled trial. *Bmj* 332, 266–270 (2006).
- [47] Camacho M, Cortal V, Abdullatif J, Zaghi S, Ruoff CM, Capasso R et al.: Myofunctional therapy to treat obstructive sleep apnea: a systematic review and meta-analysis. *Sleep* 38, 669–675 (2015).
- [48] Guimaraes KC, Drager LF, Genta PR, Marcondes BF, Lorenzi-Filho G: Effects of oropharyngeal exercises on patients with moderate obstructive sleep apnea syndrome. *Am J Respir Crit Care Med* 179, 962–966 (2009).
- [49] Randerath WJ, Galetke W, Domanski U, Weitkunat R, Ruhle KH: Tongue-muscle training by intraoral electrical neurostimulation in patients with obstructive sleep apnea. *Sleep* 27, 254–259 (2004).
- [50] Camacho M, Cortal V, Brietzke SE, Holty JE, Guilleminault C, Capasso R: Tracheostomy as treatment for adult obstructive sleep apnea: a systematic review and meta-analysis. *Laryngoscope* 124, 803–811 (2014).
- [51] Browaldh N, Nerfeldt P, Lysdahl M, Bring J, Friberg D: SKUP3 randomised controlled trial: polysomnographic results after uvulopalatopharyngoplasty in selected patients with obstructive sleep apnoea. *Thorax* 68, 846–853 (2013).
- [52] Li HY, Wang PC, Lee LA, Chen NH, Fang TJ: Prediction of uvulopalatopharyngoplasty outcome: anatomy-based staging system versus severity-based staging system. *Sleep* 29, 1537–1541 (2006).
- [53] Martí S, Sampol G, Muñoz X, Torres F, Roca A, Lloberes P et al.: Mortality in severe sleep apnoea/hypopnoea syndrome patients: impact of treatment. *Eur Respir J* 20, 1511–1518 (2002).
- [54] Goktas O, Solmaz M, Goktas G, Olze H: Long-term results in obstructive sleep apnea syndrome (OSAS) after laser-assisted uvulopalatoplasty (LAUP). *PLoS One* 9, e100211 (2014).
- [55] Sundaram S, Bridgman SA, Lim J, Lasserson TJ: Surgery for obstructive sleep apnoea. *Cochrane Database Syst Rev*, Cd001004 (2005).
- [56] Won CH, Li KK, Guilleminault C: Surgical treatment of obstructive sleep apnea: upper airway and maxillomandibular surgery. *Proc Am Thorac Soc* 5, 193–199 (2008).
- [57] Choi JH, Kim SN, Cho JH: Efficacy of the Pillar implant in the treatment of snoring and mild-to-moderate obstructive sleep apnea: a meta-analysis. *Laryngoscope* 123, 269–276 (2013).
- [58] Maurer JT, Sommer JU, Hein G, Hormann K, Heiser C, Stuck BA: Palatal implants in the treatment of obstructive sleep apnea: a randomised, placebo-controlled single-centre trial. *Eur Arch Otorhinolaryngol* 269, 1851–1856 (2012).

- [59] Back LJ, Hytonen ML, Roine RP, Malmivaara AO: Radiofrequency ablation treatment of soft palate for patients with snoring: a systematic review of effectiveness and adverse effects. *Laryngoscope* 119, 1241–1250 (2009).
- [60] Back LJ, Liukko T, Rantanen I, Peltola JS, Partinen M, Ylikoski J et al.: Radiofrequency surgery of the soft palate in the treatment of mild obstructive sleep apnea is not effective as a single-stage procedure: A randomized single-blinded placebo-controlled trial. *Laryngoscope* 119, 1621–1627 (2009).
- [61] Farrar J, Ryan J, Oliver E, Gillespie MB: Radiofrequency ablation for the treatment of obstructive sleep apnea: a meta-analysis. *Laryngoscope* 118, 1878–1883 (2008).
- [62] Strollo PJ, Jr., Soose RJ, Maurer JT, de Vries N, Cornelius J, Froymovich O et al.: Upper-airway stimulation for obstructive sleep apnea. *N Engl J Med* 370, 139–149 (2014).
- [63] Woodson BT, Gillespie MB, Soose RJ, Maurer JT, de Vries N, Steward DL et al.: Randomized controlled withdrawal study of upper airway stimulation on OSA: short- and long-term effect. *Otolaryngol Head Neck Surg* 151, 880–887 (2014).
- [64] Bangiyev JN, Traboulsi H, Abdulhamid I, Rozzelle A, Thottam PJ: Sleep architecture in Pierre-Robin sequence: The effect of mandibular distraction osteogenesis. *Int J Pediatr Otorhinolaryngol* 89, 72–75 (2016).
- [65] Vezina JP, Blumen M, Buchet I, Chabolle F: Sleep-disordered breathing: choosing the right cephalometric analysis. *J Oral Maxillofac Surg* 70, 1442–1448 (2012).
- [66] Pirklbauer K, Russmueller G, Stiebellehner L, Nell C, Sinko K, Millesi G et al.: Maxillomandibular advancement for treatment of obstructive sleep apnea syndrome: a systematic review. *J Oral Maxillofac Surg* 69, e165–176 (2011).
- [67] Holty JE, Guilleminault C: Maxillomandibular advancement for the treatment of obstructive sleep apnea: a systematic review and meta-analysis. *Sleep Med Rev* 14, 287–297 (2010).
- [68] Vicini C, Dallan I, Campanini A, De Vito A, Barbanti F, Giorgiomarrano G et al.: Surgery vs ventilation in adult severe obstructive sleep apnea syndrome. *Am J Otolaryngol* 31, 14–20 (2010).